



RESEARCH ARTICLE

Effectiveness of Polyethylene Glycol 3350 versus Lactulose in Management of Functional Constipation in Children

Hanin Ahmed Mansour, MD^{1*}, Ali Ibrahim² and Ali Mohammad³

¹Department of Pediatrics, Faculty of Medicine, Tishreen University Hospital, Lattakia, Syria

²Professor of Gastroenterology and Hepatic Diseases, Department of Pediatrics, Faculty of Medicine, Tishreen University Hospital, Lattakia, Syria

³Professor of Nutrition and Metabolism, Department of Pediatrics, Faculty of medicine, Tishreen University Hospital, Lattakia, Syria

*Corresponding author: Hanin Ahmed Mansour, MD, Department of Pediatrics, Faculty of medicine, Tishreen University Hospital, Lattakia, Syria



Abstract

Introduction: Constipation is one of the common problems in childhood. Recently, Polyethylene Glycol (PEG 3350) has been suggested as a good alternative laxative to Lactulose as a treatment option in pediatric constipation. The current study aimed to compare the efficacy of two laxatives (PEG 3350 and Lactulose) in the management of functional constipation and evaluate the side effects.

Methods and materials: In an open-label clinical trial, 1-13 year-old children with functional constipation according to ROME IV criteria and the presence of RAIR based on anorectal manometry were studied. The patients were randomized to receive either PEG3350 (0.8 g/kg/d) or Lactulose (2 ml/kg/d). Both groups received the medication orally in two divided doses for 12 weeks. They were evaluated clinically at the end of 4, 8, and 12 weeks of therapy, and the end of 16 and 20 weeks from enrolment. At the end of the 12th week, Success was defined as a defecation frequency ≥ 3 /week and improvement in stool consistency without painful defecation, as well as encopresis ≤ 1 every two weeks.

Results: We enrolled 43 patients (M20, F23) aged 4.16 \pm 2.3 years. At the end of the 12th week, good clinical outcomes were achieved in 95% (PEG) and 77.3% (Lactulose). The PEG group had a significant increase in defecations in a week compared with the Lactulose group (6.26 \pm 0.5 vs. 4.94 \pm 0.8, $p = 0.0001$) and a significant decrease in encopresis (35% vs. 10, $p = 0.01$). PEG achieved defecation without pain (0% vs. 22.7%) and less hard stools (5% vs. 18.2%, $p = 0.1$). Patients reported less abdominal pain and bloating than children using Lactulose (25% and 0% vs. 68.2% and 27.3%, respectively). We reported a significant relapse because of sudden cessation of treatment in the Lactulose group (13.6% vs. 5%, $p = 0.04$) compared with the PEG group.

Conclusion: PEG3350 compared with Lactulose provided a higher success rate, less relapse rate, and fewer side effects in the treatment of constipated children.

Keywords

Functional constipation, PEG3350, Lactulose, Laxatives

Introduction

Chronic constipation is a serious common gastrointestinal disorder in children [1]. It is very important to exclude organic causes before making the diagnosis of functional constipation, which is defined by fulfillment of at least 2 items of the ROME IV criteria [2]. Treatment includes a fiber-rich diet, parent's education, toilet training, and pharmacotherapy [3-5]. Osmotic laxatives are recommended to use as the first-choice therapy for functional constipation in children [6,7] Polyethylene Glycol (PEG) is a water-soluble, non-absorbed polymer. It has an osmotic effect on the intestinal membrane due to its high water-binding capacity [8] which leads to fluid retention and softening of stools, so it does not need to metabolize by the colonic flora [9]. It is used to clean colons before diagnostic gastrointestinal procedures in children. Lactulose is a synthetic disaccharide that is fermented by colonic bacteria, which leads to decreased colonic PH, an increase in fecal volume, in addition, to increasing colonic transit time. When it is used for a long period, it leads to a change in bacterial colonic flora which reduces

its efficacy [10]. There have been recommendations to use low doses of PEG as a choice in the treatment of functional constipated children [11,12]. The current study aimed to compare the clinical efficacy and safety of PEG 3350 and Lactulose in the treatment of childhood functional constipation.

Methods and Patients

An open-label clinical trial was performed between March 2020 and September 2021, in Tishreen University Hospital in Lattakia. We enrolled children between 1 and 13-years-old who had functional constipation according to ROME IV criteria, newly diagnosed as well as children who were previously treated ineffectively. Exclusion criteria were organic causes of constipation as Hirschprung's disease, hypothyroidism, celiac disease, neuromuscular disorders, structural gastrointestinal tract anomaly, and previous surgery of digestive tract.

Study design

During the first visit, we recorded the patient's data and did a physical examination with an evaluation of stool retention. Then many tests were performed in addition to anorectal manometry. 19 patients were excluded, 9 patients whose RAIR was absent, 3 patients had neurologic disorders, 4 patients had anal anomalies, 2 patients had celiac disease and one patient had hypothyroidism. The Parents were asked to sign an acceptance for participation in the study. After fecal disimpaction, 43 patients were randomized to receive either PEG 3350 (0.8 g/kg/d) or Lactulose (2 ml/kg/d), orally in two divided doses for 12 weeks, and then we reduced the medication gradually to avoid sudden stopping. We evaluated the patients clinically at the end of the 4th, 8th, 12th weeks of therapy and followed-up at the end of the 16th and 20th weeks of enrolment. Success was defined at the end of the 12th week by a defecation frequency ≥ 3 /week and improvement in stool consistency without painful defecation, as well as encopresis ≤ 1 every two weeks.

Patient data were analyzed by IBM SPSS statistics version 20. A p-value of 0.05 was considered statistically significant.

Results

In the current study, there were Forty-three patients (M20, F23) mean aged 4.16 ± 2.3 years. The mean duration of constipation was 11.02 ± 7.8 months. The mean number of bowel movements per week was 2.09 ± 0.6 . 81.1% had hard stools and 95.3% had painful defecations and 23.3% had encopresis. 21 patients received PEG and 22 children received Lactulose. One patient dropped out from the study from the PEG group in the 8th week. The efficacy of the treatments at weeks 4 and 8 are shown in Table 1.

Efficacy at week 12

At the end of the 12th week, we noted good clinical outcomes in 95% (PEG) and 77.3% (Lactulose) which was statistically significant (p-value 0.03). The PEG group had a significant increase in stool number per week compared with the Lactulose group, (6.26 ± 0.5 vs. 4.94 ± 0.8 , $p = 0.0001$) and a significant decrease in encopresis (35% vs. 10%, $p = 0.01$). PEG achieved less hard stools (5% vs. 18.2%, $p = 0.1$) and defecation without pain (0% vs. 22.7%). Follow-up at 16 weeks and 20 weeks were as in Table 2.

We noted a more relapse in the Lactulose group in patients who stopped receiving the medication without gradually reducing the dose compared with the PEG group (5% vs. 13.6%, $p = 0.04$) (Figure 1).

We reported significantly more side effects in patients who received Lactulose compared with the patients who received PEG. The most common adverse effects were abdominal pain and bloating (68.2% and 27.3% vs. 25% and 0% respectively). No serious adverse effects were recorded (Figure 2).

Table 1: The efficacy of the treatments at 4 and 8 weeks.

	PEG		Lactulose		p-value	
	4w	8w	4w	8w	4w	8w
Number of defecation/week (n \pm SD)	3.23 \pm 0.5	3.76 \pm 0.6	3.46 \pm 0.7	4.40 \pm 0.7	0.03	0.001
Hard stool (%)	19%	0%	45.4%	4.5%	0.06	0.3
Painful defecation (%)	42.9%	15%	63.6%	9.1%	0.1	0.5
Improvement in encopresis (%)	0%	0%	0%	0%	---	---

Table 2: Follow-up at 16 and 20 weeks.

	PEG		Lactulose		p-value	
	16w	20w	16w	20w	16w	20w
Number of defecation/week (n \pm SD)	5.46 \pm 0.9	4.46 \pm 0.5	4.29 \pm 1.1	3.58 \pm 0.9	0.002	0.003
Hard stool (%)	5%	0%	22.7%	17.6%	0.03	0.08
Painful defecation (%)	0%	0%	22.7%	17.6%	---	---
Improvement in encopresis (%)	30%	40%	4.5%	5.8%	0.01	0.004




Figure 1: Treatments efficacy and relapse.




Figure 2: Side effects of the treatments during the study.

Discussion

In the current study, 43 constipated patients were treated. Despite both medications resulting in improving the symptoms, the number of patients who were treated successfully was higher in the PEG group, which had a significantly better effect in increasing defecation frequency in a week and decreasing encopresis, in addition, to decreasing hard stools and painful defecation. A relapse was reported in patients who had stopped receiving the medication without reducing the dose, significantly higher in the Lactulose group.

1-7.5% of visits to pediatric clinics and up to 25% of referrals to pediatric gastroenterologist clinics are affected with constipation [7]. The management of functional constipation depends on clinical experience and a half to two-thirds of children with functional constipation has a successful outcome with laxative therapy for approximately 6-12 months [13].

Early and rapid stopping of medication has an important role in recurrence [13]. PEG is used routinely in the long-term treatment of constipated adults. It is shown to be more effective than placebo or Lactulose with fewer side effects [14,15]. Recently, the ESPGHAN

and NASPGHAN guidelines recommend PEG 3350 as primary therapy in children who suffer from constipation, and Lactulose can be used if PEG is unavailable [7]. Many trials compared the efficacy of PEG 3350 and Lactulose in pediatric constipation. Jarzebicka, et al. showed that PEG had more defecation per week (7.9 ± 0.6 vs. 5.7 ± 0.5) and fewer hard stools (7% vs. 13%) but it did not evaluate its effect on encopresis, because there were doubts whether parents properly reported in the child's diary [16]. Saneian, et al. reported that PEG had more than three defecations per week, after one month when compared with Lactulose and Magnesium Hydroxide [17]. Candy, et al. showed that PEG is more effective than Lactulose in increasing the number of defecations per week (9.4 ± 4.56 vs. 5.9 ± 4.29) and in preventing the recurrence of fecal impaction [18].

Voskuijl, et al. compared PEG with Lactulose in the treatment of childhood constipation. Successful treatment was achieved in 56% compared with 29% in the Lactulose group. It achieved a significant increase in defecation frequency and a decrease in encopresis [19]. Gremse DA study found that the stool frequency and form were similar for each laxative [20].

The current study showed that side effects were higher in the Lactulose group. Abdominal pain and bloating were the most common compared with PEG. Mechanisms of colonic fermentation, production of flatus, as well as osmotic effect, and increase in colonic motility account for adverse events when using Lactulose [10,21]. We reported more diarrhea in the PEG group, we think that the additional osmotic effect due to electrolytes especially sodium sulfate played a role in increase diarrhea. Our work confirms the results of the previous studies.

Conclusion

PEG 3350 is more effective than Lactulose. It has a higher success rate and less relapse rate, in addition to its fewer side effects in the treatment of constipated children.

Acknowledgments

We wish to thank Leen Doya, a doctor in the pediatric department, Aras Abdo, and Victor Al-Khoury, doctors in the pediatric surgery department for their contribution and technical assistance in the research.

Conflict of Interest Statement

None declared.

Data Availability

We can't share patient data due to our hospital's privacy policy, which is concerned with maintaining patient confidentiality and refuses to publish or share data. Also, the informed consent signed by the parents to participate in the study prevents the sharing of information with the non-study researchers.

Statement of Ethics

All parents whose children were studied gave informed consent for the sharing of this research. Ethical clearance for this study was obtained from the Ethical Committee of the University of Tishreen Hospital.

Funding Sources

None.

Author Contributions

The Authors developed and carried out a sample collection. Literature review, the data analysis and read of the final data were done by authors.

References

1. Van den Berg MM, Benninga MA, Di Lorenzo C (2006) Epidemiology of childhood constipation: A systemic review. *Am J Gastroenterol* 101: 2401-2409.
2. Hyams JS, Di Lorenzo C, Saps M, Shulman RJ, Staiano A, et al. (2016) Functional disorders: Children and adolescents. *Gastroenterology* 150: 1456-1468.
3. Jarzebicka D, Sieczkowska J, Dadalski M, Kierkus J, Ryzko J, et al. (2016) Evaluation of the effectiveness of biofeedback therapy for functional constipation in children. *Turk J Gastroenterol* 27: 433-438.
4. Bharucha AE (2000) Treatment of the severe and intractable constipation. *Curr Treat Options Gastroenterol* 7: 291-298.
5. Rao SS, Welcher KD, Pelsang RE (1997) Effects of biofeedback therapy on anorectal function in obstructive defecation. *Dig Dis Sci* 42: 2197-2205.
6. Bharucha AE, Pemberton JH, Locke GR (2013) American Gastroenterological Association technical review on constipation. *Gastroenterology* 144: 218-238.
7. Tabbers MM, DiLorenzo C, Berger MY, Faure C, Langendam MW, et al. (2014) Evaluation and treatment of functional constipation in infants and children: Evidence-based recommendations from ESPGHAN and NASPGHAN. *J Pediatr Gastroenterol Nutr* 58: 258-274.
8. Schiller LR, Emmett M, Santa Ana CA, Fordtran JS (1988) Osmotic effects of polyethylene glycol. *Gastroenterology* 94: 933-941.
9. Bekkali NLH, Hoekman DR, Liem O, Bongers MEJ, van Wijk MP, et al. (2018) Polyethylene glycol 3350 with electrolytes versus polyethylene glycol 4000 for constipation: A randomized, controlled trial. *J Pediatr Gastroenterol Nutr* 66: 10-15.
10. Candelli M, Nista EC, Zocco MA, Gasbarrini A (2001) Idiopathic chronic constipation: Pathophysiology, diagnosis and treatment. *Hepatogastroenterology* 48: 1050-1057.
11. Loening-Baucke V (2002) Polyethylene glycol without electrolytes for children with constipation and encopresis. *J Pediatr Gastroenterol Nutr* 34: 372-377.
12. Pashankar DS, Bishop WP (2001) Efficacy and optimal dose of daily polyethylene glycol 3350 for treatment of constipation and encopresis in children. *J Pediatr* 139: 428-432.
13. Poddar U (2016) Approach to constipation in children. *Indian Pediatr* 53: 319-327.
14. DiPalma JA, DeRidder PH, Orlando RC, Kolts BE, Cleveland

- MB (2000) A randomised, placebo-controlled, multicentre study of the safety and efficacy of a new polyethylene glycol laxative. *Am J Gastroenterol* 95: 446-450.
15. Corazziari E, Badiali D, Habib FL, Reboa G, Pitto G, et al. (1996) Small volume isosmotic polyethylene glycol electrolytes balanced solution (PMF-100) in treatment of chronic non-organic constipation. *Dig Dis Sci* 41: 1636-1642.
16. Jarzebicka D, Sieczkowska-Golub J, Kierkus J, Czukowski P, Kowalczyk-Kryston M, et al. (2019) PEG 3350 versus lactulose for treatment of functional constipation in children: Randomized study. *J Pediatr Gastroenterol Nutr* 68: 318-324.
17. Saneian H, Mostofizadeh N (2012) Comparing the efficacy of polyethylene glycol (PEG), magnesium hydroxide and lactulose in treatment of functional constipation in children. *Journal of Research in Medical Science* 17: S145-S149.
18. Candy DC, Edwards D, Geraint M (2006) Treatment of faecal impaction with polyethylene glycol plus electrolytes (PEG +E) followed by a double blind comparison of PEG + E versus lactulose as maintenance therapy. *J Pediatr Gastroenterol Nutr* 43: 65-70.
19. Voskuil W, de Lorijn F, Verwijs W, Hogeman P, Heijmans J, et al. (2004) PEG3350 (Transipeg) versus lactulose in the treatment of childhood functional constipation: A double, randomized, controlled multicentre trial. *Gut* 53: 1590-1594.
20. Gremse DA, Hixon J, Crutchfield A (2002) Comparison of polyethylene glycol 3350 and lactulose for treatment of chronic constipation in children. *Clin Pediatr* 41: 225-229.
21. Youssef NN, Peters JM, Henderson W, Shultz-Peters S, Lockhart DK, et al. (2002) Dose response of PEG 3350 for the treatment of childhood faecal impaction. *J Pediatr* 141: 410-414.