



REVIEW ARTICLE

Cochlear Implantation in Older Adults: Effectiveness and Expanded Indications

Nicholas J Thompson, MD*, Margaret E Richter, BA, Margaret T Dillon, AuD, and Harold C Pillsbury, MD



Department of Otolaryngology-Head & Neck Surgery, The University of North Carolina at Chapel Hill, North Carolina, USA

*Corresponding author: Nicholas J Thompson, MD, Resident Physician, Department of Otolaryngology/Head & Neck Surgery, University of North Carolina at Chapel Hill, 170 Manning Drive, CB#7070, Houtp Building, Rm G190A, NC 27599-7070, USA, Tel: 919-966-3344, Fax: 919-966-7941

Abstract

Older adults who qualify for cochlear implantation demonstrate significant improvements in speech recognition and quality of life with cochlear implant use. Cochlear implantation is a safe and effective procedure for patients with hearing loss, with some additional medical considerations specific to older adult patients. The indications for cochlear implantation have expanded from cases of bilateral severe-to-profound hearing loss to cases of normal-to-moderate low-frequency hearing sensitivity and severe-to-profound high-frequency hearing loss. The expansion of indications for cochlear implantation was prompted by evidence of low-frequency hearing preservation due to modified electrode array designs and surgical techniques, and significantly improved performance when listening with the combination of acoustic and electric stimulation (EAS) in the same ear. Older adults who underwent cochlear implantation as part of clinical trials investigating the effectiveness of EAS have been shown to experience significantly improved speech recognition and quality of life as compared to preoperative performance with conventional hearing aids.

Overview

Cochlear implantation is the standard treatment for cases of moderate-to-profound sensorineural hearing loss who experience limited speech recognition with appropriately-fit hearing aids. Though originally indicated for patients with severe-to-profound hearing loss bilaterally, candidacy criteria for cochlear implantation have expanded over the past several decades. Populations including those with normal-to-moderate low-frequency hearing in the ear-to-be implanted and those with normal hearing in the contralateral ear (known as

single-sided deafness) have demonstrated significantly improved performance with cochlear implant (CI) use as part of clinical trials, prompting the expansion of candidacy criteria to include these patient populations [1-6]. The prevalence of hearing loss increases with age, which negatively impacts quality of life and has shown to be associated with cognitive impairment [7-9]. Providing older adults (≥ 65 years of age) with effective treatment options such as cochlear implantation for their hearing loss supports improved speech recognition and quality of life [10-20]; however, insurance coverage of this procedure is currently restricted to beneficiaries with bilateral moderate-to-profound sensorineural hearing loss. The aim of the present report is to review aspects of cochlear implantation unique to older adults and data that demonstrate the effectiveness of CI use for older adults who meet the new criteria for cochlear implantation.

Medical Considerations

The overall health of an older adult patient is an important consideration during the candidacy evaluation for cochlear implantation. A combination of comorbidities, medications, and physiologic changes associated with aging can contribute to increased risk for undergoing anesthesia [21]. Specific attention should be focused on cardiovascular, pulmonary, neurological, and renal function in older adults being evaluated for preoperative clearance [22]. The overall risks of anesthesia and the surgical procedure itself should be weighed against the expected benefits from CI use.



Citation: Thompson NJ, Richter ME, Dillon MT, Pillsbury HC (2020) Cochlear Implantation in Older Adults: Effectiveness and Expanded Indications. J Geriatr Med Gerontol 6:098. doi.org/10.23937/2469-5858/1510098

Accepted: September 28, 2020; **Published:** September 30, 2020

Copyright: © 2020 Thompson NJ, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Generally, there are few systemic effects following cochlear implantation, though there are potential postoperative considerations for older adults including flap-related issues, dizziness and vertigo, and psychological considerations [23]. Flap-related issues are considered the most common complication following cochlear implantation. It is possible that chronic vascular changes from comorbidities such as long-standing hypertension, diabetes mellitus, or other vascular disease may increase this risk. Postoperative dizziness is rare following cochlear implantation, though older adults have an increased likelihood of underlying multisensory balance issues that may contribute to transient dizziness postoperatively [23]. The patient's psychological health, including attention to symptoms of dementia, should be included as part of the cochlear implantation candidacy evaluation as these may limit the overall success of CI use, though some studies have suggested CI use may slow cognitive decline [9,20,24].

Several studies examining perioperative and postoperative outcomes of older adults undergoing cochlear implantation report overall low complication rates with no long-term morbidity or mortality [11,25,26]. Though Coehlo, et al. (2009) found that advanced age was not an independent risk factor for general anesthesia, a study by Carlson, et al. (2010) showed older adults were more likely to experience anesthetic complications but found no difference in surgical complications as compared to younger adults. Cochlear implantation is generally considered to be a safe procedure for older adults that are deemed suitable to undergo general anesthesia.

Recently, some CI centers have developed methods for cochlear implantation under local anesthesia [27-30]. A modified surgical approach is used in combination with local anesthesia and minimal sedation. Preliminary studies have suggested that this can be a safe and effective alternative for patients that may not be able to tolerate general anesthesia. Studies have been performed comparing cochlear implantation under conscious sedation/local anesthesia and those undergoing general anesthesia. There reportedly are decreased surgical times, lengths of stay, and drug costs in patients implanted with conscious sedation/local anesthesia, with no differences in perioperative morbidity compared to patients implanted under general anesthesia [29,30]. Though further research and surgical training may be required before widespread use, this surgical technique may be an alternative for older adult CI candidates with multiple comorbidities that may be poor candidates for general anesthesia.

Postoperative Performance: Cochlear Implants

Speech recognition

A primary measure of postoperative success with CI use is speech recognition, which assesses the listener's understanding of spoken words. Typically, recorded word or sentence materials are presented to the listen-

er in a soundbooth to assess preoperative performance with appropriately-fit hearing aids and postoperative performance with the CI. Multiple studies demonstrate significant improvement in speech recognition scores for older adults listening with a CI as compared to their preoperative performance with hearing aids [10-15,19]. Typically, patients demonstrate significant improvements in speech recognition within the initial months of CI use and experience asymptotic performance between 6-12 months of listening experience [31]. The findings are mixed when comparing the speech recognition of younger and older adult CI recipients, with some reporting similar performance between the age groups [10-12,19] and others reporting younger adults experience better performance than older adults [13,15]. A consideration of previous comparisons between age groups is that older adults may require longer CI use than younger adults before experiencing asymptotic speech recognition performance. Dillon, et al. (2013) observed that older adults continued to experience significant improvements in speech recognition in quiet and in noise out to 5 years of CI use, and maintained superior speech recognition through 10 years. While the rate of performance growth and acclimatization for speech recognition may differ from younger adult CI recipients, an important consideration is older adults experience a significant improvement with CI use as compared to their preoperative performance with hearing aids [10-15,19].

Quality of life

Though speech recognition measures are valuable clinical tools to measure performance growth with CI use, the influence on the patient's quality of life is another important measure of the success of cochlear implantation. Older adult CI recipients report significant improvements in their overall quality of life as well as significant improvements on measures specific to their social life, confidence, emotional wellbeing, telephone use, and levels of depression [15-20]. These subjective benefits have largely been found to be similar to younger adult CI recipients, though a study by Olze, et al. (2012) suggesting older adult CI recipients may experience greater improvements in quality of life as compared to younger adult CI recipients [19]. This was speculated to be secondary to the more severe effects of hearing loss on functioning capabilities and decline of health with age of elderly individuals as well as the realistic expectations of those undergoing cochlear implantation. Collectively these studies substantiate that restoring the hearing of older adults via cochlear implantation and CI use can significantly improve quality of life and communication abilities in this population.

Postoperative Performance: Electric-Acoustic Stimulation

Electric-Acoustic Stimulation (EAS)

Indications for cochlear implantation have expanded




Figure 1: Audiogram displaying the unaided detection thresholds candidacy criteria (in gray) for the MED-EL Corporation Cochlear Implant System.

to include patients with normal-to-moderate low-to-mid frequency hearing sensitivity, severe-to-profound high-frequency hearing loss, and limited speech recognition with appropriately-fit hearing aids (Figure 1). Modified electrode array designs and surgical techniques have been shown to support low-to-mid frequency acoustic hearing preservation in the implanted ear following cochlear implantation [1,2,32-34]. In cases of postoperative hearing preservation, patients are fit with a device that combines acoustic amplification and CI technology in the same ear, known as electric-acoustic stimulation (EAS). Recipients with postoperative hearing preservation demonstrate significantly better speech recognition in quiet and noise when listening with EAS as compared to listening with a CI alone [1,2,33,35-37]. Additionally, listeners of EAS devices have shown significant improvements in localization (identifying the location of a sound source), and quality of life as compared to preoperative abilities with hearing aids [1,2,33,35,37-40]. Successful low-frequency hearing preservation has been reported for older adult CI recipients [40], supporting those older adults with hearing configurations meeting the expand-

ed indications criteria should be considered candidates for cochlear implantation.

Speech recognition

The addition of acoustic low-frequency hearing with electric stimulation provides the patient with cues that are typically not well replicated by a CI. Low-frequency cues help to resolve the fundamental frequency [37,41] and interaural timing differences [36,38,39], which support better speech recognition in noise and in challenging listening situations. Auditory aging has been shown to reduce the temporal acuity [42]. The initial clinical trials investigating the effectiveness of EAS included subjects above 65 years of age at implantation. Results are mixed regarding whether advanced age at implantation influences performance with EAS, with one study reporting a higher age at implantation yielded lower speech recognition scores [2] while another found no significant effect of age at implantation on performance [1]. A consideration of these studies is the study endpoints were 12 months post-activation. Considering older adult CI users have shown significant improvements

out to 5 years of listening experience [14], older adult EAS listeners may also experience further improvements with long-term listening experience. While it is currently unclear to the extent that age at implantation may have on speech recognition with EAS, patients - regardless of age at implantation - demonstrate a benefit with EAS over preoperative abilities with hearing aids. A similar but more general conclusion has been made by Sprinzl and Riechelmann (2010), discussing that EAS devices can be an effective treatment for the older population, but more research on the topic is needed [43].

Quality of life

As observed in CI-alone users, EAS users have reported significant improvements in quality of life as compared to their preoperative perceptions [1,2]. Subjects listening with EAS report a decrease in perceived disability and an increase in device satisfaction as compared to their preoperative listening condition [1,2]. For instance, when analyzing the results from the Abbreviated Profile of Hearing Aid Benefit (APHAB), Pillsbury, et al. (2018) found that the overall score decreased (improved) by 30.2% at the 12-month interval. The APHAB has a total score as well as scores for four subcategories, which include Ease of Communication, Reverberation, Effectiveness in Background Noise, and Aversiveness of Sound. When analyzing performance on the individual subscales, 90% of subjects reported their EAS device provided benefit for the Ease of Communication and Reverberation subscales, and 91.7% of subjects reported benefit for the Effectiveness in Background Noise subscale [1]. Gantz, et al. (2016) reported similar findings, observing significant improvements on the Ease of Communication, Reverberation, and Effectiveness in Background Noise subscales. A significant difference was not observed on the Aversiveness subscale, which is a positive outcome as the Aversiveness subscale measures whether environmental sounds are bothersome or uncomfortably loud [2]. While investigations are ongoing, it appears that older adult EAS users may perceive postoperative improvements that support an improved quality of life.

Future Directions

The candidacy criteria for cochlear implantation continue to expand with evidence from prospective clinical trials demonstrating the effectiveness of CI use in new patient populations. Recently, the FDA approved single-sided deafness and asymmetric hearing loss as indications for cochlear implantation for the MED-EL CI system. Investigations of the effectiveness of CI use in subjects with single-sided deafness or asymmetric hearing loss included older adult recipients. While prospective clinical trials demonstrate the effectiveness of CI use on measures of quality of life, localization, and speech recognition in quiet and noise for subjects with single-sided deafness and asymmetric hearing loss [5,6,44], the current criteria for Medicare beneficiaries are restricted to

those with moderate-to-profound hearing loss in both ears [45]. Ongoing work is investigating outcomes of CI use specific to older adult recipients with single-sided deafness and asymmetric hearing loss.

Conclusions

Older adult CI recipients demonstrate significant improvements in speech recognition and quality of life. Patients report significant improvements in their social life, confidence, emotional wellbeing, telephone use, and levels of depression. Indications for cochlear implantation continue to expand and now include those with single-sided deafness and asymmetric hearing loss. Older adult patients who meet the new indications for cochlear implantation may experience significantly improved speech recognition and quality of life as compared to their abilities with alternative treatment options.

Acknowledgements

Figure 1 was provided by MED-EL Corporation (Innsbruck, Austria).

Funding Source/Conflicts of Interest

MER and MTD are supported by a research grant from MED-EL Corporation to the university. HCP is a consultant for MED-EL Corporation. NJT has no conflicts of interest.

References

1. Harold C Pillsbury 3rd, Margaret T Dillon, Craig A Buchman, Hinrich Staecker, Sandra M Prentiss, et al. (2018) Multicenter us clinical trial with an Electric-Acoustic Stimulation (EAS) system in adults: Final outcomes. *Otol Neurotol* 39: 299-305.
2. Bruce J Gantz, Camille Dunn, Jacob Oleson, Marlan Hansen, Aaron Parkinson, et al. (2016) Multicenter clinical trial of the nucleus hybrid s8 cochlear implant: Final outcomes. *Laryngoscope* 126: 962-973.
3. Margaret T Dillon, Emily Buss, Meredith L Anderson, English R King, Ellen J Deres, et al. (2017) cochlear implantation in cases of unilateral hearing loss: Initial localization abilities. *Ear Hear* 38: 611-619.
4. Margaret T Dillon, Emily Buss, Meredith A Rooth, English R King, Ellen J Deres, et al. (2017) Effect of cochlear implantation on quality of life in adults with unilateral hearing loss. *Audiol Neurootol* 22: 259-271.
5. Emily Buss, Margaret T Dillon, Meredith A Rooth, English R King, Ellen J Deres, et al. (2018) Effects of cochlear implantation on binaural hearing in adults with unilateral hearing loss. *Trends Hear* 22: 2331216518771173.
6. Nicholas J Thompson, Margaret T Dillon, Emily Buss, Meredith A Rooth, English R King, et al. (2020) Subjective benefits of bimodal listening in cochlear implant recipients with asymmetric hearing loss. *Otolaryngol Head Neck Surg* 162: 933-941.
7. CD Mulrow, C Aguilar, JE Endicott, R Velez, MR Tuley, et al. (1990) Association between hearing impairment and the quality of life of elderly individuals. *J Am Geriatr Soc* 38: 45-50.

8. Frank R Lin, E Jeffrey Metter, Richard J O'Brien, Susan M Resnick, Alan B Zonderman, et al. (2011) Hearing loss and incident dementia. *Arch Neurol* 68: 214-220.
9. Alessandro Castiglione, Alice Benatti, Carmelita Velardita, Diego Favaro, Elisa Padoan, et al. (2016) Aging, cognitive decline and hearing loss: Effects of auditory rehabilitation and training with hearing aids and cochlear implants on cognitive function and depression among older adults. *Audiol Neurootol* 21: 21-28.
10. Katharina Rohloff, Mario Koopmann, Daniel Wei, Claudia Rudack, Eleftherios Savvas (2017) Cochlear implantation in the elderly: Does age matter? *Otol Neurotol* 38: 54-59.
11. Matthew L Carlson, Joseph T Breen, Rene H Gifford, Colin LW Driscoll, Brian A Neff, et al. (2010) Cochlear implantation in the octogenarian and nonagenarian. *Otol Neurotol* 31: 1343-1349.
12. RF Labadie, VN Carrasco, CH Gilmer, HC Pillsbury 3rd (2000) Cochlear implant performance in senior citizens. *Otolaryngol Head Neck Surg* 123: 419-424.
13. Veronique Chatelin, Eugene J Kim, Colin Driscoll, Jannine Larky, Colleen Polite, et al. (2004) Cochlear implant outcomes in the elderly. *Otol Neurotol* 25: 298-301.
14. Margaret T Dillon, Emily Buss, Marcia C Adunka, English R King, Harold C Pillsbury 3rd, et al. (2013) Long-term speech perception in elderly cochlear implant users. *JAMA Otolaryngol Head Neck Surg* 139: 279-283.
15. Katrien Vermeire, Jan PL Brokx, Floris L Wuyts, Ellen Cochet, Anouk Hofkens, et al. (2005) Quality-of-life benefit from cochlear implantation in the elderly. *Otol Neurotol* 26: 188-195.
16. Hamid R Djalilian, Timothy A King, Sharon L Smith, Samuel C Levine (2002) Cochlear implantation in the elderly: Results and quality-of-life assessment. *Ann Otol Rhinol Laryngol* 111: 890-895.
17. KL Horn, NB McMahon, DC McMahon, JS Lewis, M Barker, et al. (1991) Functional use of the Nucleus 22-channel cochlear implant in the elderly. *Laryngoscope* 101: 284-288.
18. Marie-Hortense Sonnet, Bettina Montaut-Verient, Jean-Yves Niemier, Michel Hoen, Laurence Ribeyre, et al. (2017) Cognitive abilities and quality of life after cochlear implantation in the elderly. *Otol Neurotol* 38: e296-e301.
19. Heidi Olze, Stefan Gräbel, Ulrike Förster, Nina Zirke, Laura E Huhnd, et al. (2012) Elderly patients benefit from cochlear implantation regarding auditory rehabilitation, quality of life, tinnitus, and stress. *Laryngoscope* 122: 196-203.
20. Isabelle Mosnier, Jean-Pierre Bebear, Mathieu Marx, Bernard Frayssé, Eric Truy, et al. (2015) Improvement of cognitive function after cochlear implantation in elderly patients. *JAMA Otolaryngol Head Neck Surg* 141: 442-450.
21. Strom C, Rasmussen LS (2014) Challenges in anaesthesia for elderly. *Singapore Dent J* 35C: 23-29.
22. Hirsch CH (1995) When your patient needs surgery: Weighing risks versus benefits. *Geriatrics* 50: 26-31.
23. Buchman CA, Fucci MJ, Luxford WM (1999) Cochlear implants in the geriatric population: Benefits outweigh risks. *Ear Nose Throat J* 78: 489-494.
24. Isabelle Mosnier, Antoine Vanier, Damien Bonnard, Geneviève Lina-Granade, Eric Truy, et al. (2018) long-term cognitive prognosis of profoundly deaf older adults after hearing rehabilitation using cochlear implants. *J Am Geriatr Soc* 66: 1553-1561.
25. Cosetti MK, Lalwani AK (2015) Is cochlear implantation safe and effective in the elderly? *Laryngoscope* 125: 1279-1281.
26. Daniel H Coelho, Joseph Yeh, Jung T Kim, Anil K Lalwani (2009) Cochlear implantation is associated with minimal anesthetic risk in the elderly. *Laryngoscope* 119: 355-358.
27. Nóra Kecskeméti, Magdolna Szőnyi, Marianna Küstel, Anita Gáborján, László Tamás, et al. (2019) Cochlear implantation under local anesthesia: A possible alternative for elderly patients. *Eur Arch Otorhinolaryngol* 276: 1643-1647.
28. Aarno Dietz, Marion Wüstefeld, Minna Niskanen, Heikki Löppönen (2016) Cochlear implant surgery in the elderly: the feasibility of a modified suprameatal approach under local anesthesia. *Otol Neurotol* 37: 487-491.
29. Samion Shabashev, Yasser Fouad, T Kate Huncke, J Thomas Roland (2017) Cochlear implantation under conscious sedation with local anesthesia; Safety, Efficacy, Costs, and Satisfaction. *Cochlear Implants Int* 18: 297-303.
30. Rogério Hamerschmidt, Ana Tereza Ramos Moreira, Gis-laine Richter Minhoto Wiemes, Sérgio Bernardo Tenório, Elizabeth Milla Tâmbara (2013) Cochlear implant surgery with local anesthesia and sedation: Comparison with general anesthesia. *Otol Neurotol* 34: 75-78.
31. Minoo Lenarz, Hasibe Sönmez, Gert Joseph, Andreas Büchner, Thomas Lenarz (2012) Long-term performance of cochlear implants in postlingually deafened adults. *Otolaryngol Head Neck Surg* 147: 112-118.
32. Turner CW, Reiss LA, Gantz BJ (2008) Combined acoustic and electric hearing: Preserving residual acoustic hearing. *Hear Res* 242: 164-171.
33. Oliver F Adunka, Margaret T Dillon, Marcia C Adunka, English R King, Harold C Pillsbury, et al. (2013) Hearing preservation and speech perception outcomes with electric-acoustic stimulation after 12 months of listening experience. *Laryngoscope* 123: 2509-2515.
34. Silke Helbig, Youssef Adel, Tobias Rader, Timo Stöver, Uwe Baumann (2016) Long-term Hearing Preservation Outcomes After Cochlear Implantation for Electric-Acoustic Stimulation. *Otol Neurotol* 37: e353- e359.
35. Margaret T Dillon, Emily Buss, Oliver F Adunka, Craig A Buchman, Harold C Pillsbury (2015) Influence of Test Condition on Speech Perception with Electric-Acoustic Stimulation. *Am J Audiol* 24: 520-528.
36. René H Gifford, Michael F Dorman, Henryk Skarzynski, Artur Lorens, Marek Polak, et al. (2013) Cochlear implantation with hearing preservation yields significant benefit for speech recognition in complex listening environments. *Ear Hear* 34: 413-425.
37. Dorman MF, Gifford RH (2010) Combining acoustic and electric stimulation in the service of speech recognition. *Int J Audiol* 49: 912-919.
38. René H Gifford, Timothy J Davis, Linsey W Sunderhaus, Christine Menapace, Barbara Buck, et al. (2017) Combined Electric and Acoustic Stimulation With Hearing Preservation: Effect of Cochlear Implant Low-Frequency Cutoff on Speech Understanding and Perceived Listening Difficulty. *Ear Hear* 38: 539-553.
39. René H Gifford, D Wesley Grantham, Sterling W Sheffield, Timothy J Davis, Robert Dwyer, et al. (2014) Localization and interaural time difference (ITD) thresholds for cochlear implant recipients with preserved acoustic hearing in the implanted ear. *Hear Res* 312: 28-37.

40. Yao WN, Turner CW, Gantz BJ (2006) Stability of low-frequency residual hearing in patients who are candidates for combined acoustic plus electric hearing. *J Speech Lang Hear Res* 49: 1085-1090.
41. Carroll J, Tiaden S, Zeng FG (2011) Fundamental frequency is critical to speech perception in noise in combined acoustic and electric hearing. *J Acoust Soc Am* 130: 2054-2062.
42. Schneider B, Pichora-Fuller K, Daneman M (2010) Effects of senescent changes in audition and cognition on spoken language comprehension. *The Aging Auditory System* 2010: 167-210.
43. Sprinzi GM, Riechelmann H (2010) Current trends in treating hearing loss in elderly people: a review of the technology and treatment options - a mini-review. *Gerontology* 56: 351-358.
44. Jill B Firszt, Ruth M Reeder, Laura K Holden, Noël Y Dwyer, Asymmetric Hearing Study Team (2018) Results in Adult Cochlear Implant Recipients With Varied Asymmetric Hearing: A Prospective Longitudinal Study of Speech Recognition, Localization, and Participant Report. *Ear Hear* 39: 845-862.
45. Steve E Phurrough, Louis B Jacques, Madeline M Ulrich, Francina Spencer, Jackie Sheridan-Moore (2015) Decision Memo for Cochlear Implantation (CAG-00107N). Centers for Medicare & Medicaid Services.