



REVIEW ARTICLE

Synovial Macrophages: Potential Key Modulators of Cartilage Damage, Osteophyte Formation and Pain in Knee Osteoarthritis

Majoska HM Berkelaar¹, Noline M Korthagen^{2,3}, Gerrit Jansen^{1#*} and Willem Evert van Spil^{4#}

¹Amsterdam Rheumatology and Immunology Center, VU University Medical Center, Amsterdam, The Netherlands

²Equine Sciences Department, Faculty of Veterinary Sciences, Utrecht University, Utrecht, The Netherlands

³Department of Orthopedics, University Medical Center Utrecht, Utrecht, The Netherlands

⁴Department of Rheumatology & Clinical Immunology, University Medical Centre Utrecht, Utrecht, The Netherlands

#Equal contribution as senior authors.

*Corresponding author: Gerrit Jansen, PhD, Amsterdam Rheumatology and Immunology Center Location VU University Medical Center, Rm CCA 2.46, De Boelelaan 1117, 1081 HV Amsterdam, The Netherlands, Tel: +31-20-444-66-85, E-mail: g.jansen@vumc.nl



Abstract

Synovitis is common in knee OA patients and a known contributor to disease incidence and progression. Macrophages are the most common immune cell type present in this inflamed synovial tissue and expectedly contribute both directly and indirectly to OA progression through the induction of inflammatory mediators, growth factors and proteinases, resulting in enhanced cartilage degeneration and osteophyte formation. Furthermore, macrophage infiltration and soluble macrophage products may be associated with pain in OA, although data are somewhat controversial. OA synovium features macrophage heterogeneity, represented by the presence of subsets covering the broad spectrum of M1 (pro-inflammatory) to M2 (anti-inflammatory) phenotypes, whose emergence corresponds to the cytokine profile found in OA patients. Conflicting results have been reported in both early and advanced OA concerning macrophage number, location, marker expression and cytokine profile, which may also be a consequence of the lack of universal definitions of disease stages and OA phenotypes. In this review, we summarize and discuss the positioning of synovial macrophages in knee OA in the context of pathogenesis, synovitis, and disease monitoring and as a target for therapeutic interventions.

Keywords

Macrophage, Osteoarthritis, Knee, Synovitis

Abbreviations

ACL: Anterior Cruciate Ligament Transaction; BMP: Bone Morphogenic Protein; CCL: CC-chemokine Ligand; CXCL: CXC-Chemokine Ligand; COX: Cyclooxygenase; DAMP: Damage-Associated Molecular Pattern; DMM: Destabilized Medial Meniscus; DMOAD: Disease-Modifying Osteoarthritic Drug; ECM: Extracellular Matrix; FPR: Formyl Peptide Receptor; FR: Folate Receptor; GM-CSF: Granulocyte Macrophage Colony-Stimulating Factor; IFN- γ : Interferon γ ; IL: Interleukin; iNOS: Inducible Nitric Oxide Synthase; KL grade: Kellgren-Lawrence Radiographic Grade of OA; LPS: Lipopolysaccharide; M-CSF: Macrophage Stimulating Factor; MIA: Monosodium Iodoacetate; MIF: Macrophage Migration Inhibitory Factor; MMP: Matrix Metalloproteinase; MRI: Magnetic Resonance Imaging; MTX: Methotrexate; NF- κ B: Nuclear Factor Kappa-Light-Chain-Enhancer of Activated B cells; NGF: Nerve Growth Factor; NO: Nitric Oxide; NSAIDs: Nonsteroidal Anti-Inflammatory Drugs; OA: Osteoarthritis; PET: Positron Emission Tomography; RA: Rheumatoid Arthritis; RANKL: Receptor Activator of NF- κ B ligand; ROS: Reactive Oxygen Species; SPECT: Single-Photon Emission Computed Tomography; TA: Triamcinolone Acetonide; Th: T Helper Cell; TGF: Tumor Growth Factor; TLR: Toll-Like Receptor; TNF: Tumor Necrosis Factor; US: Ultrasonography

Introduction

Osteoarthritis (OA) is among the ten leading causes of disability in the Western world [1-3], affecting the elderly as well as individuals of working age [4]. OA is generally a slowly progressive joint disease and is characterized by pain, stiffness, and sometimes swelling of the joint [5,6], all of which can result in impaired function [1,3]. Current non-surgical treatment options focus on symptom alleviation, including analgesics, anti-inflammatory therapy and physical and occupational therapy. However, these treatments do not suffice as they do not modify the course of the underlying disease [7,8]. Hence, due to the high economic and personal burden related to OA, there is a large and growing demand for disease-modifying therapies (DMOADs).

To date, the pathogenesis of OA has not been elucidated completely. Previously, OA was merely considered a disease of 'wear and tear' [9-12] with cartilage degradation as its key characteristic [3]. Nowadays, OA is considered a multifactorial disease affecting the whole organ (Figure 1), resulting in structural and func-

tional changes in many articular and periarticular tissues [2,9,10,13]. Moreover, OA is considered a heterogeneous disease, leading to a variety of disease manifestations between patients, joints, and disease stages [2,9,10,13,14]. Due to this heterogeneity, it is hypothesized that stratification of OA subtypes is warranted in order to provide appropriate treatments.

Synovitis is a common feature in OA, but its severity and timing differ among patients [2,5,10,12,13,15-18]. Synovitis does seem to correspond to pain [10]. Furthermore, several studies have demonstrated an association between synovitis and (radiographic) OA progression [2,5,9,16] in general and the advancement of cartilage breakdown in particular [12,18,19]. Therefore, treating synovitis in OA would potentially be beneficial, as both symptom alleviation and reduction of joint degradation could be achieved [12]. To this end, DMOADs may serve as effective treatments of synovitis [12,16].

A possible treatment strategy for OA could be the modulation of infiltrating cells. In rheumatic joints, neutrophils and lymphocytes generally comprise the ma-




Figure 1: Schematic overview of a healthy knee joint (left) and osteoarthritic knee joint (right). The cardinal features of OA are cartilage damage, osteophyte formation and subchondral sclerosis, and synovitis. Synovial macrophages play a critical role in cartilage degeneration and osteophyte formation by the induction of pro-inflammatory cytokines (e.g. TNF- α and IL-1 β), growth factors and matrix metalloproteinases (MMPs), resulting in clinical symptoms of OA. Abbreviations: IL-1 β , interleukin-1 β ; TNF- α , tumor necrosis factor α .

majority of leukocytes and this is reflected in the synovial fluid in active rheumatoid arthritis (RA) [20]. In contrast, macrophages are relatively more present in the synovial fluid in OA as compared to RA [21]. Furthermore, macrophages are also found to be the predominant inflammatory cell type in the synovial tissue in OA [12,22,23]. Their role in OA, however, is less well understood.

Macrophages represent a dynamic cell type harboring the ability to alter their pro- or anti-inflammatory phenotype and function upon environmental stimuli [24]. Several studies have demonstrated that synovial macrophage infiltration is positively correlated with OA progression and disease severity [25,26]. Moreover, synovial macrophages have been associated with cartilage degradation [27], osteophyte formation [28], and pain [29] in OA (Figure 1). As macrophages are key cells in the pathogenesis of OA, modulating synovial macrophages might be sufficient to alleviate OA symptoms and prevent progression.

The potential role of synovial macrophages in knee OA will be further discussed in the following paragraphs of this review. To ensure a comprehensive overview a literature (Supplementary File) search strategy was employed as described in Supplementary Figure 1.

The synovium in healthy and OA joints

The synovial membrane constitutes a boundary between the joint and surrounding musculoskeletal tissues [30]. Through the secretion of synovial fluid, synovium conserves articular mobility and limits friction between articular cartilage surfaces [6,31]. Additionally, the synovial fluid nourishes the articular cartilage and removes metabolites and other matrix degradation products from the joint [12].

The synovium consists of two layers: the synovial lining or intima, comprising 2 to 3 cell layers of synovial macrophages (type A synoviocytes) and fibroblasts (type B synoviocytes); and the synovial sublining or subintima, containing collagen type III, blood vessels and a few immune cells, including macrophages [6,30-32]. Synovial fibroblasts primarily produce synovial fluid components, including hyaluron [30,31], whereas synovial macrophages remove joint debris from the synovial fluid, clear bacterial infections, and control the pro-inflammatory and anti-inflammatory cytokine balance in the synovial fluid [30]. Therefore, both cell types are vital for the maintenance of joint homeostasis.

Clinical monitoring of inflamed OA synovium

Multiple methodologies are available for detecting and grading of synovitis in OA and other joint diseases, including arthroscopy, synovial biopsy, and non-invasive imaging techniques such as magnetic resonance imaging (MRI) and ultrasonography (US) [16]. Synovitis is detected in 50% of the patients using arthroscopy [18] and in 47% and 73% of the patients using US and MRI,

respectively [12]. Typically, synovitis cannot be visualized by X-ray imaging.

Arthroscopy portrays synovitis as localized proliferative alterations in the synovium with signs of inflammation, such as an increased vascularity [12]. MRI can be used to determine the presence and/or extent of synovial thickening and effusion [12,33]. US can be used to monitor synovial hypertrophy, hyperemia, and joint effusion [12,19]. The latter non-invasive techniques are increasingly used for assessing synovitis, as their findings correlate to arthroscopic findings [12,16] and/or histopathologically observed synovitis [12,33].

Histological aberrancies in the OA synovium include synovial hypertrophy and hyperplasia, accompanied by infiltration of mononuclear cells in the synovial sublining and lining layer [11,12,34]. Inflamed areas in the OA synovium show increased vascularization and cellular infiltration [18]. Generally, synovitis is found in close proximity to damaged cartilage [12,18,35], but no preferential location has been found so far [16].

Synovitis in different OA disease stages

The extent of synovitis differs between disease severity levels, disease stages, and patients [16]. In general, synovitis is present in both early and advanced OA, but tends to increase with disease severity [16,19,36,37]. Studies by Ayril, et al. demonstrated that in advanced OA, characterized by excessive cartilage damage, synovitis detected by arthroscopy was more prominent as compared to early-stage OA [18]. However, other studies have failed to detect synovitis [36,37] or showed synovitis in less than 50% of advanced OA patients [38]. Moreover, synovitis can precede the radiographic presentation of knee OA [39,40], indicating that synovitis might even predate cartilage change [39]. These findings underline the great variety in synovitis between OA patients, but at the same time its potential importance.

Synovial macrophages in OA

Synovial macrophages are located in the synovial lining and scattered throughout the synovial sublining in OA, mostly confined to sites of cartilage damage [26,37,41-43]. Even though synovial macrophage numbers are lower in OA than in RA [44], macrophages are the most prominent immune cell type [34,45-47] and are highly activated [29,42] in OA.

Animal studies by Blom, et al. suggest that macrophage activation can occur due to cartilage damage in OA [28,48]. Mediators from cartilage probably leak into the synovial fluid and activate synovial macrophages (Figure 1). Potential mediators include damage-associated molecular patterns (DAMPs), including extracellular matrix (ECM) components released by damaged cartilage, e.g. fibrinogen and plasma proteins, such as α 1m, α 2m and Gc-globulin [49], alarmins [50], and basic calcium phosphate crystals [49,51]. DAMPs predominantly

signal via Toll-like receptor (TLR) 4, expressed by synovial macrophages, and CD14, which forms a complex with TLR-2 and TLR-4 to initiate DAMP-related macrophage activation [25,52]. In OA, the expression of macrophage markers, including CD14 and MHC class II genes, is associated with joint space narrowing and osteophyte formation [26].


Activation of synovial macrophages leads to release of tumour necrosis factor (TNF)- α and other pro-inflammatory cytokines [49,51], catabolic mediators, such as matrix metalloproteinases (MMPs) [12,28], and anabolic factors, which can induce osteophyte formation [28]. The notion that release of cartilage breakdown products can trigger activation of macrophages and cytokine production indicates that synovitis is part of a vicious cycle of inflammation and cartilage breakdown [53]. In this cascade of events, the activation mechanism of macrophages in OA and their role as either initiator and/or drivers of OA remains to be defined.

Activated synovial macrophages in OA can also stimulate other immune cells, particularly T cells. In fact, the mononuclear infiltration is mainly comprised of CD4+ T cells [34,45,46]. Macrophages can take up fragments of

type II collagen in OA, present them on their surface, and activate T cells [35]. Studies by Shen, et al. show that CD4+ T cell numbers increase at OA onset in the anterior cruciate ligament transection (ACLT) model. Furthermore, the increased numbers of CD4+ T cells were followed by a rise in numbers of synovial macrophages, which can possibly be explained by the stimulation of CCL9 release by CD4+ T cells [54]. Together, these findings suggest a close interplay between macrophages and CD4+ T cells in OA.

Detecting macrophages in OA synovitis

Currently, synovial macrophages can only be visualized histologically. However, non-invasive tools for visualizing macrophages are under development. Macrophage Positron Emission Tomography (PET) imaging using the macrophage G-protein coupled receptor formyl peptide receptor (FPR)-1 tracer showed encouraging results in experimental OA [55]. Another approach utilizes Etarfolatide, which has a high binding affinity for the folate receptor (FR)- β expressed on activated macrophages [25]. Single-photon emission computed tomography (SPECT) with (99m)Tc-EC20 (Etarfolatide) has been successfully applied to detect macrophages in



knee joints of patients with varying degrees of OA [29]. Lastly, a recent study demonstrated macrophage visualization in OA using SPECT with ¹¹¹In-Octreoscan, which has a high affinity for the somatostatin receptor subtype 2 expressed by macrophages [56].

Macrophage polarization and subsets

Macrophages are dynamic cells that can respond to stimuli in their microenvironment through modification of their phenotype and function [57]. Because of this, macrophages are key players in initiation as well as resolution of inflammation [57]. Although a spectrum of macrophage subsets and even mixed subtypes of macrophages exists [58], a crude subdivision can be made in two extremes: “M1” classically activated and “M2” alternatively activated macrophages (Figure 2). M2 macrophages can be further classified in subgroups, including M2a, M2b and M2c, based on activation stimuli and functions [59-61].

M1 and M2 macrophages exert different functions, related to their distinct cytokine release profile. M1 macrophages are involved in Th1 stimulation and are characterized by IL-12, IL-23, TNF- α , IL-1 and nitric oxide (NO) release. M2 macrophages stimulate Th2 cells [42,60] and predominantly release IL-10, CCL18, TGF- β and IL-1 receptor antagonist (IL-1RA) [42,57,61] upon activation (Figure 2). In turn, M1 and M2 macrophages are activated through cytokines produced by Th1 and Th2 cells, respectively [42]. M1 activation occurs through interferon γ (IFN- γ), the TLR ligand lipopolysaccharide (LPS) and components of intracellular bacteria. Skewing to M2 macrophages proceeds upon exposure to interleukin IL-4, IL-13, parasites, complement factors, macrophage stimulating factor (M-CSF), IL-10 and tumor growth factor (TGF)- β [42,57,59] (Figure 2).

Originally, M1 macrophages were considered pro-inflammatory, while M2 macrophages were considered anti-inflammatory through production of anti-inflammatory and regulatory cytokines and chemokines [60]. This gave rise to the assumption that M2 macrophages are beneficial in inflammatory diseases and might be involved in tissue repair and wound healing [59]. However, this view has been (partially) revisited as M2 macrophages have also been associated with chronic inflammation in cancer and other diseases [59,62] and have been shown to be able to produce pro-inflammatory cytokines in a synovial microenvironment with auto-antibodies in RA patients [63,64].

In addition to their distinct function and cytokine release, M1 and M2 macrophages can be distinguished through differentially expressed receptors and other markers. CD14 [26], a marker of monocytes and macrophages, is more abundantly expressed on M2 compared to M1 macrophages [25]. CD163 [25,57,62,65,66] and CD206 [67] are considered preferred markers of M2 macrophages. M1 macrophages predominantly express MHC class II, CD68, CD80 and CD86 [66,68] (Figure 2).

Although Folate Receptor β (FR β) expression has been identified as a marker for activated macrophages in RA [69-71], its expression on polarized macrophages is incompletely known; in tumour-associated macrophages FR β is expressed on M2 macrophages [72], but in OA it is also expressed on M1 macrophages [42]. Thus, caution should be taken when classifying FR β + macrophages as M2 macrophages solely on the basis of CD163 co-expression as both receptors seem to have a mixed pattern of expression on M1 and M2 macrophages in RA and OA [42].

Macrophage polarization in knee OA

The inflammatory milieu of the synovium in OA has been implicated in the skewing of macrophages to both pro-inflammatory and anti-inflammatory phenotypes. In this respect, Utomo, et al. demonstrated that most synovial macrophages in OA have a M2 phenotype characterized by IL-10 production [62]. However, multiple studies demonstrated that both M1 and M2 macrophage phenotypes are present in OA [24,28,41,66,68,73,74]. In fact, synovial macrophages in OA were identified to actually express a combination of M1 and M2 phenotypic markers, including CD14, CD163 and CD86 [24,34,36,73,75-77] (Figure 2). It has been hypothesized that the combined expression of both M1 and M2 macrophage markers might be caused by synovial macrophages being “trapped in transition” [25,29]. The exact mechanism underlying activation of M1 and M2 synovial macrophages in OA is still unresolved.

In early stages of OA, increased skewing of bone marrow monocyte precursors towards the M1 phenotype has been reported [74]. However, macrophages stimulated with synovial fluid of early OA resulted in increased release of CCL2 and IL6 by mesenchymal stromal cells (MSC), which in turn can direct M2 polarization of macrophages [78]. Studies by Fahy, et al. report a large intra-patient variation in phenotypic marker expression patterns between OA synovium regions. Specifically, CD86 expression was either absent or present on macrophages in the synovial lining or sublining, while expression of CD206 was most pronounced on macrophages in the lining layer [67]. Likewise, Manferdini, et al. mainly identified CD80 M1 macrophages in the OA synovium lining layer, while CD206-positive M2 macrophages were predominantly located in the sublining layer [68]. Moreover, Manferdini, et al. reported a large inter-patient variability in the M1 and M2 marker expression in the OA synovium [68]. Hence, generalizations of the phenotypic marker expression patterns should be exercised with caution.

Several studies have demonstrated differences between RA and OA synovial macrophage phenotypes. Lower expression of CD14+ macrophages was found in OA as compared to RA synovium [68], whereas TGF- β expression was higher in OA [42]. Furthermore, macrophages in OA synovium had lower FR β expression than

RA macrophages, and FR β + /CD163+ macrophages were more dominant than FR β + /CD163- macrophages in the lining layer of the OA synovium [42]. Together, relatively high M2 marker expression is demonstrated on OA synovial macrophages although absolute expression appears lower than on RA synovial macrophages.

Cytokines and chemokines related to synovial macrophages

In general, higher inflammatory cytokine levels are observed in RA compared with OA synovium, although both diseases show partly analogous cytokine profiles, consisting of elevated levels of IL-1 β , TNF- α , IL-6 and IL-4 [44,79-82]. However, there are also potentially important differences, such as the granulocyte macrophage colony-stimulating factor (GM-CSF) expression that is more prominent in OA compared with RA [44]. In OA, anti-inflammatory cytokine levels, e.g. IL-10, are increased compared to healthy controls [73], which may be consistent with the suggested dominance of M2 over M1 macrophages in OA.

IL-1 β and TNF- α constitute the key mediators released by activated pro-inflammatory macrophages and staining patterns of these mediators correspond to the distribution of synovial macrophages in OA [24]. Both IL-1 β and TNF- α are primarily detected in the lining layer and to a lesser extent in the sublining layer of OA synovium [24,27,83]. Furthermore, IL-1 β and TNF- α induce pro-inflammatory cytokine production, including IL-6 and IL-8, and thereby stimulate inflammation. Moreover, IL-1 β suppresses type II collagen, which is a key constituent of cartilage, and subsequently aggravates cartilage damage [27]. Together, these findings suggest that synovial macrophages of mixed phenotypes account for the pro-inflammatory and anti-inflammatory cytokine production to fuel the pathogenesis of OA. However, Beekhuizen, et al. did not observe elevated levels of TNF- α and IL-1 β in advanced OA [84], suggesting that these cytokines may not be essential in this phase of OA.

Macrophages are not the only cell type in the joint capable of producing these indicated cytokines and the exact cellular source of remains largely unclear [24]. Furthermore, a study by Manferdini, et al. demonstrates that cell cultures containing both synovial fibroblasts and macrophages produced more pro-inflammatory mediators, including IL-6 and IL-8, compared to cell cultures with synovial fibroblasts alone [73]. This finding illustrates that macrophages are directly involved in the induction of inflammatory cytokine release from synovial fibroblasts, but mechanisms driving this process remain to be determined.

Chemokines are small secreted factors that mediate recruitment of cells [85]. Several OA studies found increased levels of chemokines involved in the attraction of macrophages. CCL2 is one of the most important

chemokines involved in macrophage attraction in chronic inflammation [73,86] and elevated levels of CCL2 promote OA development [87]. CCL13 is involved in recruitment of monocytes and T lymphocytes to joints. Gao, et al. found a positive correlation between serum and synovial fluid levels of CCL13 and radiographic severity of knee OA [88]. Additionally, CXCL12 is involved in macrophage recruitment and CXCL12 plasma and synovial fluid levels are correlated with radiographic OA disease severity, which could be explained by their role in macrophage attraction [89]. Other mediators involved in macrophage attraction include CCL22, which is elevated in the synovial fluid of OA patients [84], and CCL3, of which plasma levels are positively correlated with radiographic OA disease severity [90]. Chemokines may thus mediate the attraction of macrophages to the joint and the worsening of OA.

Synovial macrophages in different OA disease stages

Studies on CD68 expression suggest a pronounced macrophage infiltration in synovium of patients with early OA compared to advanced OA [73,91]. This coincided with an increased expression of nuclear factor (NF)- κ B, inducible nitric oxide synthase (iNOS) and other pro-inflammatory mediators by macrophages in early OA [39,93]. Ene, et al. reported high levels of inflammatory cytokines, including IL-6, IL-1, TNF- α , and a greater mononuclear cell infiltrate in early compared to advanced OA [92]. Consistently, Benito, et al. reported increased TNF- α , IL-1 β and VEGF expression and mononuclear cell infiltration in early compared with advanced OA [91]. Lastly, Ning, et al. confirmed an increase in macrophage-related mediators in early compared to advanced OA [93].

Oppositely, other studies observed similar CD68+ macrophages numbers in synovium of early and advanced OA [83,94]. Additionally, similar synovial fluid TNF- α levels were found in early and advanced OA, whereas IL-1 β expression was increased in the advanced stage OA group [83]. The reason for this apparent discrepancy in results is not clear.

It has been hypothesized that innate immunity is more prominent in early OA, while adaptive immunity dominates in advanced OA. Firstly, larger lymphoid infiltrations were observed in the synovium in advanced OA compared to early OA [38]. Secondly, the more extensive immune cell infiltration in advanced as compared to early OA could be explained by the subsequent activation of adaptive immunity. Thirdly, this hypothesis could explain the increase of mediators such as VEGF, IL-4, IL-6, IL-8, CCL3 and CCL4 in advanced stage OA and their positive correlation to radiographic severity of OA [77,80,91,95].

Macrophage-induced cartilage damage

Several preclinical and clinical studies have shown that synovitis is linked to cartilage damage in OA [96].

Macrophages are the predominant immune cells in synovitis [27] and CD14 deficiency is correlated to delayed cartilage degradation in destabilized medial meniscus (DMM) OA models in mice [53]. Utomo, et al. studied the impact of factors released by macrophage subsets on cartilage degradation in an *in vitro* cartilage explant model. Their results demonstrated that pro-inflammatory M1 products stimulate cartilage damage, whereas anti-inflammatory M2 factors failed to directly inhibit cartilage degeneration or inflammation [97]. Furthermore, a recent study has shown that M1 mediators inhibit mesenchymal stem cell chondrogenic capacity and thereby exacerbate cartilage damage [68].

Inflammatory mediators released by macrophages at least partially induce cartilage damage by both downregulating matrix anabolism by chondrocytes and inducing release of MMPs, aggrecanases and other inflammatory mediators by synovial fibroblasts [77,94,98] (Figure 1). Bondeson, et al. depleted synovial macrophages by anti-CD14 conjugated magnetic beads in synovial cell cultures derived from digested OA synovium, which resulted in reduced production of IL-1 β , TNF- α and MMPs by synovial fibroblasts and in less cartilage damage [22]. Furthermore, cartilage degradation was decreased in IL-1 β and TNF- α neutralisation experiments [22].

Macrophages induce cartilage damage through the release of MMPs, including MMP-1, -3 and -9 [48], and through the secretion of cytokines, which in turn lead to proteinase secretion by synovial fibroblasts [99]. Depletion of macrophages from OA synovium resulted in lower expression of MMP-2, -3, and -9 [22,48], as well as CCL2 [22]. CCL2 is a chemokine involved in macrophage attraction and normally associated with increased MMP-3 and MMP-13 production [87]. MMP-3 release is also increased by TNF- α and IL-1 β stimulation [100], both key mediators released by macrophages.

In addition to MMP-induced cartilage damage, activated macrophages increase cartilage degradation through the induction of oxidative stress within the joint. Studies by Steinbeck, et al. showed that in early OA patients, synovial macrophages produced reactive oxygen species (ROS), hypochlorous acid, chlorine gas and chlorinated peptides, which exert an oxidative effect on cartilage [101]. Consistently, myeloperoxidase, an enzyme responsible for the production of chlorine gas, hypochlorous acid and chlorinated peptides, appeared elevated in early OA compared to controls and end-stage OA [101]. Furthermore, a high level of iNOS, which induces nitric oxide (NO) production, has been detected in OA patients [38]. Specifically, double immunohistochemical staining showed strong iNOS expression in CD68+ macrophages in the synovial lining and sublining layer of advanced OA patients [102]. The release of NO results in oxidative stress and impacted immune processes mediating cartilage integrity [102]. These findings were confirmed in a recent study, showing that NO production in synovial macrophages was

significantly elevated in modest to severe knee OA patients compared to controls [103].

Macrophage-mediated osteophyte formation

Next to cartilage degeneration, the MMP release by activated synovial macrophages also permits osteophyte formation and/or progression through cartilage matrix remodelling [104,105]. Osteophytes are bony outgrowths at the joint margin (Figure 1) and their presence is associated with pain in OA [12]. In early collagenase-induced OA, synovitis is considered a key contributor to osteophyte formation [106]. Synovial macrophages also release growth factors that can induce osteophyte formation, such as TGF- β [28] (Figure 1). Moreover, macrophage cell surface marker levels were directly proportional to osteophyte formation [25].

A study in collagenase-induced OA murine knee joints revealed that depletion of synovial macrophages resulted in an 85% reduction of osteophyte formation after 7 days [28]. This animal model shows resemblance with human knee OA and is claimed to be a suitable model for studying osteophyte formation during OA. Remarkably, in another mouse model of papain-induced OA treated with triamcinolone acetonide (TA), Siebelt, et al. reported that enhanced macrophage infiltration diminished osteophyte formation [66]. This effect might have been caused by increased proportions of CD163/FR β -positive M2 anti-inflammatory macrophages, which, in *in vitro* cultures, displayed increased IL-10 mRNA expression [66]. These findings underline the potential importance of macrophage polarization in osteophyte formation in OA.

Osteophyte formation is regulated by growth factors produced by synovial macrophages, in particular TGF- β [28]. Van Lent, et al. showed that macrophages induce osteophyte formation through the production of TGF- β and by the expression of factors that induce chondrogenesis of mesenchymal cells [107]. Furthermore, their studies highlighted the possible crucial role of bone morphogenic proteins (BMPs) in osteophyte formation. Additionally, in patients with advanced OA, hepatocyte growth factor (HGF) stimulated the macrophage-selective production of TGF- β 1 and BMP-2 [76]. These findings pinpoint TGF- β and BMP as key players in osteophyte formation in OA, even though macrophages might not be the exclusive producers of these growth factors [28].

In addition to growth factor production, the role of the alarmin proteins S100A8/A9 in osteophyte formation has been studied in collagenase-induced OA, DMM-induced OA and in patients with early OA [104,105]. Alarmins are released by activated macrophages and signal through TLR-4 [106,108]. Elevated alarmin S100A8/A9 plasma levels in patients with early knee OA correlated with development of osteophytes over 2 and 5 years [104]. Further research is required to uncover a possible mechanistic relationship between alarmin secretion and osteophyte formation.

In OA, subchondral bone remodeling is spatially variable and related to disease stage [109]. Bone resorption is increased in early stages, whereas in advanced stages ectopic bone formation predominates [109]. Macrophages can influence this process of bone remodeling, which involves bone resorption by osteoclasts and formation by osteoblasts [109]. The multinucleated osteoclasts originate from the same precursor cells as macrophages through stimulation of M-CSF and receptor activator of NF- κ B ligand (RANKL) [110,111]. Moreover, macrophages are able to induce osteoclastogenesis through the production of inflammatory cytokines, including IL-1, TNF and IL-6 [110]. These cytokines promote osteoclastogenesis both directly by stimulating osteoclast precursors and indirectly by inducing RANKL on synovial fibroblasts [110,111].

Several studies have shown that macrophages isolated from the synovial fluid of knee OA patients differentiated into osteoclasts and showed lacunar resorption upon RANKL stimulation [112-114]. However, Adamopoulos, et al. reported greater osteoclast differentiation upon RANKL stimulation of RA synovial fluid macrophages as compared to OA synovial fluid macrophages [112,113]. Nevertheless, although osteoclast differentiation of synovial fluid macrophages may be more relevant in RA, the induction of osteoclast formation by macrophages could still be a contributing mechanism involved in, for instance, osteophyte formation in OA.

Macrophage infiltration correlates with pain

Synovitis has been recognized as one potential determinant of pain in OA patients and in animal models [2,38,77,115,116]. Synovitis as observed by MRI is related to knee pain [117], even more when using contrast-enhanced MRI [118,119] or dynamic contrast-enhanced MRI [120]. However, another study reported that changes in synovitis were not associated with concomitant changes in pain [96]. Also, in an animal model of collagenase-induced OA, the peak of synovitis did not correlate with the detection of pain [121]. Whether these contradictory findings relate to different OA phenotypes in patients and/or are merely limitations of animal OA models, warrants further investigation [116].

The specific association between macrophages and OA pain has been described in a monosodium iodoacetate (MIA) rat model of OA [116]. Furthermore, macrophage infiltration correlated with pain in patients with varying degrees of human knee OA [29]. These infiltrating macrophages were activated based on shedding of CD163 and CD14 markers and were associated with pain [29]. Another study confirmed the association of macrophage CD14 expression with pain in patients with knee OA [25]. Furthermore, elevated expression of CCL2 in synovium was found in patients with symptomatic OA and CCL2 level in the synovial fluid of these patients was associated with pain [85]. Conceivably, these mediators play a role in OA-related pain through the attraction of

macrophages and other immune cells, resulting in an augmented immune cell infiltrate [115].

Apart from macrophage infiltration, several studies have demonstrated that cytokine release by macrophages is also related to pain in OA. Notably, serum levels of TNF- α were associated with pain in patients with symptomatic OA [122], whereas in severe OA, IL-6 release correlated with pain [123]. IL-6 can be secreted by either macrophages or T cells, but the cellular source of IL-6 in OA is unknown [123]. Synovial fluid levels of macrophage migration inhibitory factor (MIF), a pro-inflammatory cytokine mainly produced by macrophages, was positively correlated with pain in a study of patients with Kellgren-Lawrence (KL) grade 2-4 knee OA [124]. MIF induces pro-inflammatory cytokine release by macrophages, which might explain this correlation. Finally, synovial macrophages displayed increased nerve growth factor (NGF) expression, which was associated with pain in OA [77]. Together, these findings underscore a positive correlation between soluble macrophage mediators and pain in knee OA.

Nevertheless, preliminary findings of a study in end-stage knee OA demonstrated that higher levels of GM-CSF and its receptor CD116 were correlated with reduced pain, independently of the expression of other macrophage markers [125]. This seems counterintuitive since GM-CSF and its receptor are involved in macrophage survival and attraction [126]. However, in early collagenase-induced OA, pain was reported to be GM-CSF dependent [126]. Given the fact that stimulation of monocytic progenitor cells might be more pronounced in early disease, this could explain the observed differences between early and advanced OA.

Studies by Klein-Wieringa, et al. and De Jong, et al. demonstrate that presence of synovial CD4+ T cells rather than macrophages is associated with knee pain in advanced OA [45,46]. Nevertheless, an indirect role of macrophages may still be envisioned as they are involved in activation of CD4+ T cells and T cells activate macrophages in return.

Synovial macrophage-related OA therapies

Effects of intra-articular therapies on macrophages: Glucocorticoids slightly reduced CD68+ macrophages in the lining layer in patients with symptomatic knee OA, but this did not translate into different MCP-1, MIP-1 alpha, MMP-1, MMP-3, TIMP-1, and TIMP-2 levels in the synovial lining and sublining layers [127]. Rochetti, et al. reported a reduction of macrophage numbers in advanced knee OA after intra-articular injection of hyaluronan or methylprednisolone [128]. Mechanistically, this involved two processes; hyaluronan appeared to mainly stimulate reparative processes, whereas the corticosteroid appeared to reduce the inflammatory process [128].

Intra-articular therapies might not only reduce mac-

rophage numbers, but also modulate macrophage phenotypes. Glucocorticoids increase synovial macrophage expression of CD163, a proposed marker of M2 macrophages producing IL-10 [25,62,65,66]. Utomo, et al. added dexamethasone to synovium explants of OA patients and showed an anti-inflammatory effect in gene expression analyses [62]. In addition, dexamethasone was added to primary human monocytes that were first polarized *in vitro* in either M1 or M2 phenotypes. Dexamethasone suppressed the pro-inflammatory M1 macrophages and enhanced the anti-inflammatory M2 macrophages. Similar experiments were performed with rapamycin, bone morphogenetic protein 7 (BMP-7) and pravastatin. Generally, rapamycin and BMP-7 enhanced the inflammatory response in synovium explants and suppressed M2 macrophages. Pravastatin had no impact on the inflammatory status of the explants but did suppress M2 macrophages. Furthermore, because of different results between untreated synovium explants and explants that were pre-treated with IFN- γ and TNF- α , it was concluded that the impact of the tested compounds may be different between disease severity levels [62].

In a rat model of papain-induced OA, intra-articular triamcinolone acetonide (TA) injections resulted in reduced osteophyte formation but did not affect cartilage degeneration or subchondral sclerosis. Mechanistically, *in vitro* experiments showed that TA elicited its effect by inducing monocyte differentiation to M2 macrophages [66].

A currently investigated approach aiming at macrophage skewing is Tissuegene-C, a cell-mediated gene therapy modality for localized delivery of TGF- β 1. In a rat MIA model, IL-10 production and other M2 macrophage markers were elevated in knee joints of the Tissuegene-C group as compared to the control group, although M1 macrophage numbers were similar. These findings suggest that Tissuegene-C induces an anti-inflammatory environment in the knee joint [129]. Tissuegene-C is currently being tested in phase II clinical trials in knee OA patients [129].

Macrophage-related cytokines as treatment target:

So far, targeting cytokines has yielded disappointing results [7]. For instance, anti-NGF- β therapy in knee OA patients resulted in substantial pain reduction but was accompanied by serious side effects [7]. Furthermore, anti-IL-1 β treatment has failed to improve clinical outcomes in OA [7], even though IL-1 is a key cytokine produced by activated macrophages. Finally, anti-TNF therapy has shown similar, disappointing results in OA [7].

The lack of response in OA patients may be a result of their heterogeneity. Schue, et al. compared high and low cellularity subgroups of radiographic knee OA with respect to response to anti-TNF treatment with infliximab. At baseline, the high cellularity subgroup showed increased numbers of mononuclear cells and blood ves-

sels and higher COX-2 and IL-1 expression compared with the low cellularity subgroup, whereas other cytokine levels were similar. Remarkably, only patients with high synovial cellular infiltrate responded to anti-TNF treatment although groups were small [130]. This finding highlights the heterogeneity between patients and its potential importance for the response to treatment.

Moderate, positive effects were observed by broadly-acting anti-inflammatory drugs. For example, in ACLT rabbits, statins inhibited CCL2 and MMPs, reduced infiltration of CD68+ macrophages in the OA subintima [99] and decreased articular cartilage degradation. Likewise, CCL9 neutralization reduced macrophage and CD4+ T cell infiltration and pro-inflammatory IL-1 β expression in ACLT mice, which translated into less severe histopathological signs of OA, decreased osteoclast formation, and decreased MMP-13 expression [131]. Furthermore, a study in patients with advanced knee OA revealed that celecoxib treatment decreased macrophage infiltration and cytokine expression in the synovial membrane, independent of cyclooxygenase (COX)-2 inhibition [132].

Methotrexate (MTX) may exert a positive effect on OA as the folate receptor β on synovial macrophages may serve as an entry route [71]. A pragmatic phase III trial of MTX in knee OA patients is currently running [133]. However, given the lower number of FR β positive macrophages in OA as compared to RA, its efficacy might be less in OA [42]. To this end, assessment of FR β expression by non-invasive macrophage imaging [25,29] could prove a useful diagnostic tool to identify OA patients eligible for MTX therapy.

Summary, conclusions and future directions

Synovial macrophages are thought to be key players in OA and appear to play a role in the characteristic osteophyte formation, cartilage degeneration and pain. Several studies have suggested that macrophages in OA synovium are the main producers of inflammatory mediators, which in turn induce release of proteinases, aggrecans and growth factors by synovial fibroblasts. Although activators of synovial macrophages remain unknown, it is hypothesized that ECM debris activates macrophages in OA, which in turn aggravates the anabolic and catabolic imbalance in the OA joint. Therefore, synovial macrophages in synovitis are considered as critical drivers of OA.

In synovitis, synovial macrophages are mostly present in the lining layer of the synovium and irregularly distributed in the synovial sublining layer. These macrophages display both M1 and M2 phenotypes, as evidenced by expression of cell surface markers, gene expression, and cytokine profiles. Although the M1/M2 dichotomy is a highly simplified concept and mainly based on *in vitro* observations, it is a helpful tool. Unfortunately, in most OA studies, macrophage polarization is not fully characterized. Nevertheless, it has become clear

that the polarization state of synovial macrophages varies between OA patients and between disease severity stages. More research is warranted to gain insight in the role of distinct macrophage phenotypes.

The role of macrophages might differ in early and advanced OA. It is hypothesized that innate immunity plays a key role in early OA pathology, whereas in advanced stages adaptive immunity is more prominent. However, macrophages attribute to both immune responses. Macrophages have mainly been studied in animal models, ranging between DMM, ACLT, collagenase-induced and spontaneous OA models. However, differences in models should be noted and cautious interpretation of results is warranted. The surgically-induced DMM model of OA is not associated with inflammation, although minor inflammation can be triggered by the surgical procedure, whereas the collagenase-induced model is associated with significant synovitis [50]. As a result, different models may draw different conclusions about the role of macrophages in OA.

To conclude, as synovial macrophages are considered key drivers of synovitis and are associated with OA progression, targeting synovial macrophages represents a potential treatment approach. Therapies involved in macrophage phenotype modulation are promising; however, more knowledge is required about macrophage polarization in different disease stages and between OA phenotypes. Synovial macrophage imaging and marker characterization might be able to identify knee OA patients eligible for macrophage-targeted therapies. Hence, future research should focus on the role of macrophages in early and advanced OA and between different phenotypes of OA patients.

References

- Cross M, Smith E, Hoy D, Nolte S, Ackerman I, et al. (2014) The global burden of hip and knee osteoarthritis: Estimates from the global burden of disease 2010 study. *Ann Rheum Dis* 73: 1323-1330.
- Loeser RF, Goldring SR, Scanzello CR, Goldring MB (2012) Osteoarthritis: A disease of the joint as an organ. *Arthritis Rheum* 64: 1697-1707.
- Cucchiari M, de Girolamo L, Filardo G, Oliveira JM, Patrick Orth, et al. (2016) Basic science of osteoarthritis. *J Exp Orthop* 3: 22.
- Bondeson J, Blom AB, Wainwright S, Hughes C, Caterson B, et al. (2010) The role of synovial macrophages and macrophage-produced mediators in driving inflammatory and destructive responses in osteoarthritis. *Arthritis Rheum* 62: 647-657.
- Pelletier JP, Martel-Pelletier J, Abramson SB (2001) Osteoarthritis, an inflammatory disease: Potential implication for the selection of new therapeutic targets. *Arthritis Rheum* 44: 1237-1247.
- Hügler T, Geurts J (2016) What drives osteoarthritis?—Synovial versus subchondral bone pathology. *Rheumatology* 56: 1461-1471.
- Chevalier X, Eymard F, Richette P (2013) Biologic agents in osteoarthritis: Hopes and disappointments. *Nat Rev Rheumatol* 9: 400-410.
- Hochberg MC, Altman RD, April KT, Benkhalti M, Guyatt G, et al. (2012) American College of Rheumatology 2012 recommendations for the use of nonpharmacologic and pharmacologic therapies in osteoarthritis of the hand, hip, and knee. *Arthritis Care Res (Hoboken)* 64: 465-474.
- Mobasheri A, Batt M (2016) An update on the pathophysiology of osteoarthritis. *Ann Phys Rehabil Med* 59: 333-339.
- Bijlsma JW, Berenbaum F, Lafeber FP (2011) Osteoarthritis: An update with relevance for clinical practice. *Lancet* 377: 2115-2126.
- Samuels J, Krasnokutsky S, Abramson SB (2008) Osteoarthritis: A tale of three tissues. *Bull NYU Hosp Jt Dis* 66: 244-250.
- Sellam J, Berenbaum F (2010) The role of synovitis in pathophysiology and clinical symptoms of osteoarthritis. *Nat Rev Rheumatol* 6: 625-635.
- Berenbaum F (2013) Osteoarthritis as an inflammatory disease (osteoarthritis is not osteoarthrosis!). *Osteoarthritis Cartilage* 21: 16-21.
- Kraus VB, Blanco FJ, Englund M, Karsdal MA, Lohmander LS (2015) Call for standardized definitions of osteoarthritis and risk stratification for clinical trials and clinical use. *Osteoarthritis Cartilage* 23: 1233-1241.
- Wallace G, Cro S, Doré C, King L, Kluzek S, et al. (2016) Associations between clinical evidence of inflammation and synovitis in symptomatic knee osteoarthritis: A substudy of the VIDEO trial. *Arthritis Care Res* 69: 1340-1348.
- Scanzello CR, Goldring SR (2012) The role of synovitis in osteoarthritis pathogenesis. *Bone* 51: 249-257.
- Scanzello CR (2017) Role of low-grade inflammation in osteoarthritis. *Curr Opin Rheumatol* 29: 79-85.
- Ayral X, Pickering EH, Woodworth TG, Mackillop N, Dougados M (2005) Synovitis: A potential predictive factor of structural progression of medial tibiofemoral knee osteoarthritis—Results of a 1 year longitudinal arthroscopic study in 422 patients. *Osteoarthritis Cartilage* 13: 361-367.
- D'Agostino MA, Conaghan P, Le Bars M, Baron G, Grassi W, et al. (2005) EULAR report on the use of ultrasonography in painful knee osteoarthritis. Part 1: Prevalence of inflammation in osteoarthritis. *Ann Rheum Dis* 64: 1703-1709.
- Edwards SW, Hallett MB (1997) Seeing the wood for the trees: The forgotten role of neutrophils in rheumatoid arthritis. *Immunol Today* 18: 320-324.
- Fort JG, Flanigan M, Smith JB (1995) Mononuclear cell (MNC) subtypes in osteoarthritis synovial fluid. Comparison with MNC subtypes in rheumatoid arthritis synovial fluid. *J Rheumatol* 22: 1335-1337.
- Bondeson J, Wainwright SD, Lauder S, Amos N, Hughes CE (2006) The role of synovial macrophages and macrophage-produced cytokines in driving aggrecanases, matrix metalloproteinases, and other destructive and inflammatory responses in osteoarthritis. *Arthritis Res Ther* 8: 187.
- Hamilton JA, Tak PP (2009) The dynamics of macrophage lineage populations in inflammatory and autoimmune diseases. *Arthritis Rheum* 60: 1210-1221.
- de Lange-Brokaar BJE, Ioan-Facsinay A, van Osch GJVM, Zuurmond AM, Schoones J, et al. (2012) Synovial inflammation, immune cells and their cytokines in osteoarthritis: A review. *Osteoarthritis Cartilage* 20: 1484-1499.
- Daghestani HN, Pieper CF, Kraus VB (2015) Soluble macrophage biomarkers indicate inflammatory phenotypes in

- patients with knee osteoarthritis. *Arthritis Rheumatol* 67: 956-965.
26. Blom AB, Van Lent P, Van Den Bosch M, Cats H, Van Den Hoogen F, et al. (2014) Identification of synovial genes and pathways associated with disease progression in a cohort of early osteoarthritis patients (CHECK). *Osteoarthritis and Cartilage* 22: S23-S24.
27. Mabey T, Honsawek S (2015) Cytokines as biochemical markers for knee osteoarthritis. *World J Orthop* 6: 95-105.
28. Blom AB, van Lent PL, Holthuysen AE, van der Kraan PM, Roth J, et al. (2004) Synovial lining macrophages mediate osteophyte formation during experimental osteoarthritis. *Osteoarthritis Cartilage* 12: 627-635.
29. Kraus VB, McDaniel G, Huebner JL, Stabler TV, Pieper CF, et al. (2016) Direct in vivo evidence of activated macrophages in human osteoarthritis. *Osteoarthritis Cartilage* 24: 1613-1621.
30. Bhattaram P, Chandrasekharan U (2017) The joint synovium: A critical determinant of articular cartilage fate in inflammatory joint diseases. *Semin Cell Dev Biol* 62: 86-93.
31. Smith MD (2011) The normal synovium. *Open Rheumatol J* 5: 100-106.
32. Iwanaga T, Shikichi M, Kitamura H, Yanase H, Nozawa-Inoue K (2000) Morphology and functional roles of synovial cells in the joint. *Arch Histol Cytol* 63: 17-31.
33. Loeuille D, Chary-Valckenaere I, Champigneulle J, Rat AC, Toussaint F, et al. (2005) Macroscopic and microscopic features of synovial membrane inflammation in the osteoarthritic knee: Correlating magnetic resonance imaging findings with disease severity. *Arthritis Rheum* 52: 3492-3501.
34. Siebuhr A, Bay-Jensen A, Jordan J, Kjelgaard-Petersen C, Christiansen C, et al. (2016) Inflammation (or synovitis)-driven osteoarthritis: An opportunity for personalizing prognosis and treatment? *Scand J Rheumatol* 45: 87-98.
35. Saito I, Koshino T, Nakashima K, Uesugi M, Saito T (2002) Increased cellular infiltrate in inflammatory synovia of osteoarthritic knees. *Osteoarthritis Cartilage* 10: 156-162.
36. Oehler S, Neureiter D, Meyer-Scholten C, Aigner T (2002) Subtyping of osteoarthritic synoviopathy. *Clin Exp Rheumatol* 20: 633-640.
37. Deligne C, Casulli S, Pigenet A, Bougault C, Campillo-Gimenez L, et al. (2015) Differential expression of interleukin-17 and interleukin-22 in inflamed and non-inflamed synovium from osteoarthritis patients. *Osteoarthritis Cartilage* 23: 1843-1852.
38. Wyatt LA, Moreton BJ, Mapp PI, Wilson D, Hill R, et al. (2017) Histopathological subgroups in knee osteoarthritis. *Osteoarthritis Cartilage* 25: 14-22.
39. Ostojic M, Soljic V, Vukojevic K, Dapic T (2017) Immunohistochemical characterization of early and advanced knee osteoarthritis by NF- κ B and iNOS expression. *J Orthop Res* 35: 1990-1997.
40. Kema I, Kisand K, Suutre S, Murde M, Tamm A, et al. (2014) The ADAM12 is upregulated in synovitis and postinflammatory fibrosis of the synovial membrane in patients with early radiographic osteoarthritis. *Joint Bone Spine* 81: 51-56.
41. Favero M, Ramonda R, Goldring MB, Goldring SR, Punzi L (2015) Early knee osteoarthritis. *RMD Open* 1: e000062.
42. Ashraf S, Mapp PI, Walsh DA (2011) Contributions of angiogenesis to inflammation, joint damage, and pain in a rat model of osteoarthritis. *Arthritis Rheum* 63: 2700-2710.
43. Tsuneyoshi Y, Tanaka M, Nagai T, Sunahara N, Matsuda T, et al. (2012) Functional folate receptor beta-expressing macrophages in osteoarthritis synovium and their M1/M2 expression profiles. *Scand J Rheumatol* 41: 132-140.
44. Hogg N, Palmer DG, Revell PA (1985) Mononuclear phagocytes of normal and rheumatoid synovial membrane identified by monoclonal antibodies. *Immunology* 56: 673-681.
45. Farahat MN, Yanni G, Poston R, Panayi GS (1993) Cytokine expression in synovial membranes of patients with rheumatoid arthritis and osteoarthritis. *Ann Rheum Dis* 52: 870-875.
46. Klein-Wieringa IR, de Lange-Brokaar BJ, Yusuf E, Andersen SN, Kwekkeboom JC, et al. (2016) Inflammatory cells in patients with endstage knee osteoarthritis: A comparison between the synovium and the infrapatellar fat pad. *J Rheumatol* 43: 771-778.
47. De Jong AJ, De Lange-Brokaar BJ, Klein-Wieringa IR, Heijink M, Hoekstra A, et al. (2016) Synovial CD4+T cells associate with pain in osteoarthritis: Is there a role for fatty acids? *Osteoarthr Cartil* 24: S321.
48. Haywood L, McWilliams DF, Pearson CI, Gill SE, Ganesan A, et al. (2003) Inflammation and angiogenesis in osteoarthritis. *Arthritis Rheum* 48: 2173-2177.
49. Blom AB, van Lent PL, Libregts S, Holthuysen AE, van der Kraan PM, et al. (2007) Crucial role of macrophages in matrix metalloproteinase-mediated cartilage destruction during experimental osteoarthritis: Involvement of matrix metalloproteinase 3. *Arthritis Rheum* 56: 147-157.
50. Sohn DH, Sokolove J, Sharpe O, Erhart JC, Chandra PE, et al. (2012) Plasma proteins present in osteoarthritic synovial fluid can stimulate cytokine production via Toll-like receptor 4. *Arthritis Res Ther* 14: 7.
51. Liu-Bryan R (2013) Synovium and the innate inflammatory network in osteoarthritis progression topical collection on osteoarthritis. *Curr Rheumatol Rep* 15: 323-330.
52. Corr EM, Cunningham CC, Helbert L, McCarthy GM, Dunne A (2017) Osteoarthritis-associated basic calcium phosphate crystals activate membrane proximal kinases in human innate immune cells. *Arthritis Res Ther* 19: 23.
53. Sambamurthy N, Nguyen V, Smalley R, Dodge GR, Scanzello CR (2016) CD14 deficiency delays progression of cartilage degeneration and protects against early deficits in functional outcomes in a murine osteoarthritis model. *Arthritis Rheumatol* 68: 2763-2764.
54. Moradi B, Rosshirt N, Tripel E, Kirsch J, Barie A, et al. (2015) Unicompartamental and bicompartmental knee osteoarthritis show different patterns of mononuclear cell infiltration and cytokine release in the affected joints. *Clin Exp Immunol* 180: 143-154.
55. Shen PC, Wu CL, Jou IM, Lee CH, Juan HY, et al. (2011) T helper cells promote disease progression of osteoarthritis by inducing macrophage inflammatory protein-1 γ . *Osteoarthr Cartil* 19: 728-736.
56. Yang X, Chordia MD, Du X, Graves JL, Zhang Y, et al. (2016) Targeting formyl peptide receptor 1 of activated macrophages to monitor inflammation of experimental osteoarthritis in rat. *J Orthop Res* 34: 1529-1538.
57. Van Tiel ST, Utomo L, De Swart J, De Jong M, Bernsen M, et al. (2016) Imaging inflammation in the knee joint with 111-inocteoscan. *Osteoarthr Cartil* 24: S320.
58. Udalova IA, Mantovani A, Feldmann M (2016) Macrophage heterogeneity in the context of rheumatoid arthritis. *Nat Rev Rheum* 12: 472-485.

59. Xue J, Schmidt SV, Sander J, Draffehn A, Krebs W, et al. (2014) Transcriptome-based network analysis reveals a spectrum model of human macrophage activation. *Immunity* 40: 274-288.
60. Roszer T (2015) Understanding the mysterious M2 macrophage through activation markers and effector mechanisms. *Mediators Inflamm* 2015: 1-16.
61. Wang N, Liang H, Zen K (2014) Molecular mechanisms that influence the macrophage M1-M2 polarization balance. *Front Immunol* 5: 1-9.
62. Martinez FO, Gordon S (2014) The M1 and M2 paradigm of macrophage activation: Time for reassessment. *F1000Prime Rep* 6: 13.
63. Utomo L, van Osch GJ, Bayon Y, Verhaar J, Bastiaansen-Jenniskens YM (2016) Guiding synovial inflammation by macrophage phenotype modulation: An in vitro study towards a therapy for osteoarthritis. *Osteoarthr Cartilage* 24: 1629-1638.
64. Vogelpoel LTC, Hansen IS, Rispens T, Muller FJM, van Capel TMM, et al. (2014) Fc gamma receptor-TLR cross-talk elicits pro-inflammatory cytokine production by human M2 macrophages. *Nat Commun* 5: 5444.
65. Clavel C, Ceccato L, Anquetil F, Serre G, Sebbag M (2016) Among human macrophages polarised to different phenotypes, the M-CSF-oriented cells present the highest pro-inflammatory response to the rheumatoid arthritis-specific immune complexes containing ACPA. *Ann Rheum Dis* 75: 2184-2191.
66. Daghestani HN, McDaniel G, Huebner JL, Stabler T, Pieper C, et al. (2013) Soluble macrophage biomarkers are associated with osteoarthritis-related knee inflammation. *Osteoarthr Cartil* 21: S76-S77.
67. Siebelt M, Korthagen N, Wei W, Groen H, Bastiaansen-Jenniskens Y, et al. (2015) Triamcinolone acetonide activates an anti-inflammatory and folate receptor-positive macrophage that prevents osteophytosis in vivo. *Arthritis Res Ther* 17: 352.
68. Fahy N, de Vries-van Melle ML, Lehmann J, Wei W, Grotenhuis N, et al. (2014) Human osteoarthritic synovium impacts chondrogenic differentiation of mesenchymal stem cells via macrophage polarisation state. *Osteoarthr Cartil* 22: 1167-1175.
69. Manfredini C, Paoletta F, Gabusi E, Gambari L, Piacentini A, et al. (2017) Adipose stromal cells mediated switching of the pro-inflammatory profile of M1-like macrophages is facilitated by PGE2: In vitro evaluation. *Osteoarthr Cartil* 25: 1161-1171.
70. Nakashima-Matsushita N, Homma T, Yu S, Matsuda T, Sunahara N, et al. (1999) Selective expression of folate receptor beta and its possible role in methotrexate transport in synovial macrophages from patients with rheumatoid arthritis. *Arthritis Rheum* 42: 1609-1616.
71. Xia W, Hilgenbrink AR, Matteson EL, Lockwood MB, Cheng JX, et al. (2009) A functional folate receptor is induced during macrophage activation and can be used to target drugs to activated macrophages. *Blood* 113: 438-446.
72. van der Heijden JW, Oerlemans R, Dijkmans BAC, Qi H, van der Laken CJ, et al. (2009) Folate receptor beta as a potential delivery route for novel folate antagonists to macrophages in the synovial tissue of rheumatoid arthritis patients. *Arthritis Rheum* 60: 12-21.
73. Puig-Kroger A, Sierra-Filardi E, Dominguez-Soto A, Samaniego R, Corcuera MT, et al. (2009) Folate receptor beta is expressed by tumor-associated macrophages and constitutes a marker for M2 anti-inflammatory/regulatory macrophages. *Cancer Res* 69: 9395-9403.
74. Manfredini C, Paoletta F, Gabusi E, Silvestri Y, Gambari L, et al. (2016) From osteoarthritic synovium to synovial-derived cells characterization: synovial macrophages are key effector cells. *Arthritis Res Ther* 18: 83.
75. Geven EJW, Sloetjes A, DiCeglie I, De Munter W, Van Dalen S, et al. (2015) THU0026 A systemic shift towards a pro-inflammatory monocyte subpopulation in bone marrow upon locally induced experimental osteoarthritis. *Ann Rheum Dis* 74: 202.
76. Chimenti MS, Bergamini A, Triggianese P, Baffari E, Conigliaro P, et al. (2015) AB0046 Synovial fluid from rheumatoid arthritis patients modulates the immunophenotype and viability of monocytes. *Ann Rheum Dis* 74: 15-21.
77. Dankbar B, Neugebauer K, Wunrau C, Tibesku CO, Skwara A, et al. (2007) Hepatocyte growth factor induction of macrophage chemoattractant protein-1 and osteophyte-inducing factors in osteoarthritis. *J Orthop Res* 25: 569-577.
78. Stoppiello LA, Mapp PI, Wilson D, Hill R, Scammell BE, et al. (2014) Structural associations of symptomatic knee osteoarthritis. *Arthritis Rheumatol* 66: 3018-3027.
79. Gómez-Aristizábal A, Sharma A, Bakooshi MA, Kapoor M, Gilbert PM, et al. (2017) Stage-specific differences in secretory profile of mesenchymal stromal cells (MSCs) subjected to early- vs late-stage OA synovial fluid. *Osteoarthr Cartil* 25: 737-741.
80. Radojèiae MR, Thudium CS, Henriksen K, Tan K, Karlsten R, et al. (2017) Biomarker of extracellular matrix remodeling C1M and proinflammatory cytokine IL-6 are related to synovitis and pain in endstage knee osteoarthritis patients. *Pain* 158: 1254-1263.
81. Mabey T, Honsawek S, Tanavalee A, Yuktanandana P, Wilairatana V, et al. (2016) Plasma and synovial fluid inflammatory cytokine profiles in primary knee osteoarthritis. *Biomarkers* 21: 639-644.
82. Lafeber FPJG, van Spil WE (2013) Osteoarthritis year 2013 in review: Biomarkers; reflecting before moving forward, one step at a time. *Osteoarthr Cartil* 21: 1452-1464.
83. Al-Madol MA, Shaqura M, John T, Likar R, Ebied RS, et al. (2017) Comparative Expression Analyses of Pro- versus Anti-Inflammatory Mediators within Synovium of Patients with Joint Trauma, Osteoarthritis, and Rheumatoid Arthritis. *Mediators Inflamm*.
84. Scanzello CR, Umoh E, Pessler F, Diaz-Torne C, Miles T, et al. (2009) Local cytokine profiles in knee osteoarthritis: elevated synovial fluid interleukin-15 differentiates early from end-stage disease. *Osteoarthritis Cartilage* 17: 1040-1048.
85. Beekhuizen M, Gierman LM, van Spil WE, Van Osch GJVM, Huizinga TWJ, et al. (2013) An explorative study comparing levels of soluble mediators in control and osteoarthritic synovial fluid. *Osteoarthritis Cartilage* 21: 918-922.
86. Li L, Jiang BE (2015) Serum and synovial fluid chemokine ligand 2/monocyte chemoattractant protein 1 concentrations correlates with symptomatic severity in patients with knee osteoarthritis. *Ann Clin Biochem* 52: 276-282.
87. Wei Y, Bai L (2016) Recent advances in the understanding of molecular mechanisms of cartilage degeneration, synovitis and subchondral bone changes in osteoarthritis. *Connect Tissue Res* 57: 245-261.
88. Xu Y, Ke Y, Wang B, Lin J (2015) The role of MCP-1-CCR2 ligand-receptor axis in chondrocyte degradation and disease progress in knee osteoarthritis. *Biol Res* 48: 64.

89. Gao F, Tian J, Pan H, Gao J, Yao M (2015) Association of CCL13 levels in serum and synovial fluid with the radiographic severity of knee osteoarthritis. *J Investig Med* 63: 545-547.
90. He W, Wang M, Wang Y, Wang Q, Luo B (2016) Plasma and synovial fluid CXCL12 levels are correlated with disease severity in patients with knee osteoarthritis. *J Arthroplasty* 31: 373-377.
91. Zhao XY, Yang ZB, Zhang ZJ, Zhang ZQ, Kang Y, et al. (2015) CCL3 serves as a potential plasma biomarker in knee degeneration (osteoarthritis). *Osteoarthr Cartil* 23: 1405-1411.
92. Benito MJ, Veale DJ, FitzGerald O, van den Berg WB, Bresnihan B (2005) Synovial tissue inflammation in early and late osteoarthritis. *Ann Rheum Dis* 64: 1263-1267.
93. Ene R, Diana Sinescu R, Ene P, Mihaela Cîrstoiu M, Catalin Cîrstoiu F (2015) Synovial inflammation in patients with different stages of knee osteoarthritis. *Rom J Morphol Embryol* 56: 169-173.
94. Ning L, Ishijima M, Kaneko H, Kurihara H, Arikawa-Hirasawa E, et al. (2011) Correlations between both the expression levels of inflammatory mediators and growth factor in medial perimeniscal synovial tissue and the severity of medial knee osteoarthritis. *Int Orthop* 35: 831-838.
95. Rubenhagen R, Schuttrumpf JP, Sturmer KM, Frosch KH (2012) Interleukin-7 levels in synovial fluid increase with age and MMP-1 levels decrease with progression of osteoarthritis. *Acta Orthop* 83: 59-64.
96. de Lange-Brokaar BJ, Ioan-Facsinay A, Yusuf E, Kroon HM, Zuurmond AM, et al. (2016) Evolution of synovitis in osteoarthritic knees and its association with clinical features. *Osteoarthritis Cartilage* 24: 1867-1874.
97. Utomo L, Bastiaansen-Jenniskens YM, Verhaar JA, van Osch GJVM (2016) Cartilage inflammation and degeneration is enhanced by pro-inflammatory (M1) macrophages in vitro, but not inhibited directly by anti-inflammatory (M2) macrophages. *Osteoarthritis Cartilage* 24: 2162-2170.
98. Rosshirt N, Hadrian P, Elena T, Gotterbarm T, Babak M (2016) Impact of pro-/anti-inflammatory mononuclear cell infiltration on MMP/ADAMTS concentration in osteoarthritic knee joints. *Osteoarthritis Cartilage* 24: S332-S333.
99. Akasaki Y, Matsuda S, Nakayama K, Fukagawa S, Miura H, et al. (2009) Mevastatin reduces cartilage degradation in rabbit experimental osteoarthritis through inhibition of synovial inflammation. *Osteoarthritis Cartilage* 17: 235-243.
100. Kjellaard-Petersen C, Siebuhr AS, Christiansen T, Ladel C, Karsdal M, et al. (2015) Synovitis biomarkers: Ex vivo characterization of three biomarkers for identification of inflammatory osteoarthritis. *Biomarkers* 20: 547-556.
101. Steinbeck MJ, Nesti LJ, Sharkey PF, Parvizi J (2007) Myeloperoxidase and chlorinated peptides in osteoarthritis: Potential biomarkers of the disease. *J Orthop Res* 25: 1128-1135.
102. Grabowski PS, Wright PK, Van't Hof RJ, Helfrich MH, Ohshima H, et al. (1997) Immunolocalization of inducible nitric oxide synthase in synovium and cartilage in rheumatoid arthritis and osteoarthritis. *Br J Rheumatol* 36: 651-655.
103. Suantawee T, Tantavisut S, Adisakwattana S, Tanpowpong T, Tanavalee A, et al. (2015) Upregulation of inducible nitric oxide synthase and nitrotyrosine expression in primary knee osteoarthritis. *J Med Assoc Thai* 98: S91-S97.
104. Schelbergen RF, De Munter W, Van Den Bosch MH, Lafeber FP, Sloetjes A, et al. (2016) Alarmins S100A8/S100A9 aggravate osteophyte formation in experimental osteoarthritis and predict osteophyte progression in early human symptomatic osteoarthritis. *Ann Rheum Dis* 75: 218-225.
105. Van Lent PLEM, Blom AB, Schelbergen RF, Sloetjes A, Lafeber FP, et al. (2012) Active involvement of alarmins S100A8 and S100A9 in the regulation of synovial activation and joint destruction during mouse and human osteoarthritis. *Arthritis Rheum* 64: 1466-1476.
106. Van den Bosch MH, Blom AB, Schelbergen RF, Vogl T, Roth JP, et al. (2016) Induction of Canonical Wnt Signaling by the Alarmins S100A8/A9 in Murine Knee Joints: Implications for Osteoarthritis. *Arthritis Rheumatol* 68: 152-163.
107. Van Lent PLEM, Blom AB, van der Kraan P, Holthuysen AEM, Vitters E, et al. (2004) Crucial role of synovial lining macrophages in the promotion of transforming growth factor beta-mediated osteophyte formation. *Arthritis Rheum* 50: 103-111.
108. Schelbergen RF, Blom AB, Van Den Bosch MH, Sloetjes A, Abdollahi-Roodsaz S, et al. (2012) Alarmins S100A8 and S100A9 elicit a catabolic effect in human osteoarthritic chondrocytes that is dependent on toll-like receptor 4. *Arthritis Rheum* 64: 1477-1487.
109. Burr DB, Gallant MA (2012) Bone remodelling in osteoarthritis. *Nat Rev Rheumatol* 8: 665-673.
110. Nakashima T, Takayanagi H (2009) Osteoimmunology: Crosstalk between the immune and bone systems. *J Clin Immunol* 29: 555-567.
111. Boyle WJ, Simonet WS, Lacey DL (2003) Osteoclast differentiation and activation. *Nature* 423: 337-342.
112. Adamopoulos IE, Danks L, Itonaga I, Locklin RM, Sabokbar A, et al. (2006) Stimulation of osteoclast formation by inflammatory synovial fluid. *Virchows Arch* 449: 69-77.
113. Adamopoulos IE, Sabokbar A, Wordsworth BP, Carr A, Ferguson DJ, et al. (2006) Synovial fluid macrophages are capable of osteoclast formation and resorption. *J Pathol* 208: 35-43.
114. Danks L, Sabokbar A, Gundle R, Athanasou NA (2002) Synovial macrophage-osteoclast differentiation in inflammatory arthritis. *Ann Rheum Dis* 61: 916-921.
115. Wyatt LA, Mapp PI, Wilson D, Hill R, Scammell BE, et al. (2016) Gene expression patterns in the synovium and their association with symptomatic knee osteoarthritis. *Osteoarthritis Cartilage* 24: S28-S29.
116. Nwosu LN, Mapp PI, Chapman V, Walsh DA (2016) Relationship between structural pathology and pain behaviour in a model of osteoarthritis (OA). *Osteoarthritis Cartilage* 24: 1910-1917.
117. Hill CL, Hunter DJ, Niu J, Clancy M, Guermazi A, et al. (2007) Synovitis detected on magnetic resonance imaging and its relation to pain and cartilage loss in knee osteoarthritis. *Ann Rheum Dis* 66: 1599-1603.
118. Baker K, Grainger A, Niu J, Clancy M, Guermazi A, et al. (2010) Relation of synovitis to knee pain using contrast-enhanced MRIs. *Ann Rheum Dis* 69: 1779-1783.
119. Krasnokutsky S, Belitskaya-Lévy I, Bencardino J, Samuels J, Attur M, et al. (2011) Quantitative magnetic resonance imaging evidence of synovial proliferation is associated with radiographic severity of knee osteoarthritis. *Arthritis Rheum* 63: 2983-2991.
120. Gait AD, Hodgson R, Parkes MJ, Hutchinson CE, O'Neill TW, et al. (2016) Synovial volume vs synovial measurements from dynamic contrast enhanced MRI as measures of response in osteoarthritis. *Osteoarthritis Cartilage* 24: 1392-1398.

121. Golds EE, Cooke TD, Poole AR (1983) Immune regulation of collagenase secretion in rheumatoid and osteoarthritic synovial cell cultures. *Coll Relat Res* 3: 125-140.
122. Stannus OP, Jones G, Blizzard L, Cicuttini FM, Ding C (2013) Associations between serum levels of inflammatory markers and change in knee pain over 5 years in older adults: A prospective cohort study. *Ann Rheum Dis* 72: 535-540.
123. Imamura M, Ezquerro F, Marcon Alfieri F, Vilas Boas L, Tozetto-Mendoza TR, et al. (2015) Serum levels of proinflammatory cytokines in painful knee osteoarthritis and sensitization. *Int J Inflamm* 2015: 329792.
124. Zhang PL, Liu J, Xu L, Sun Y, Sun XC (2016) Synovial fluid macrophage migration inhibitory factor levels correlate with severity of self-reported pain in knee osteoarthritis patients. *Med Sci Monit* 22: 2182-2186.
125. Pustjens MF, Mastbergen SC, Eijkelkamp N, Lafeber FP (2015) Expression of GM-CSF and its receptor CD116 in the synovium of OA patients is negatively correlated with pain. *Osteoarthr Cartil* 24: 923-924.
126. Cook AD, Pobjoy J, Steidl S, Durr M, Braine EL, et al. (2012) Granulocyte-macrophage colony-stimulating factor is a key mediator in experimental osteoarthritis pain and disease development. *Arthritis Res Ther* 14: 199.
127. Young L, Katrib A, Cuello C, Vollmer-Conna U, Bertouch JV, et al. (2001) Effects of intraarticular glucocorticoids on macrophage infiltration and mediators of joint damage in osteoarthritis synovial membranes: Findings in a double-blind, placebo-controlled study. *Arthritis Rheum* 44: 343-350.
128. Ronchetti IP, Guerra D, Taparelli F, Boraldi F, Bergamini G, et al. (2001) Morphological analysis of knee synovial membrane biopsies from a randomized controlled clinical study comparing the effects of sodium hyaluronate (Hyalgan) and methylprednisolone acetate (Depomedrol) in osteoarthritis. *Rheumatology* 40: 158-169.
129. Choi H (2016) Tissuegene-c (INVOSSA™) Induces an anti-inflammatory environment in the arthritic knee joint via macrophage polarization. *Osteoarthritis Cartilage* 24: S335.
130. Schue JR, Tawfik O, Bolce R, Smith DD, Hinson G, et al. (2012) Intraarticular infliximab for knee osteoarthritis: High baseline levels of synovial cellularity and high MRI cartilage injury at the lateral tibial plateau predict improvement in total WOMAC score. *Arthritis Rheum* 64: S485.
131. Shen PC, Lu CS, Shiau AL, Lee CH, Jou IM, et al. (2013) Lentiviral small hairpin RNA knockdown of macrophage inflammatory protein-1gamma ameliorates experimentally induced osteoarthritis in mice. *Hum Gene Ther* 24: 871-882.
132. Alvarez-Soria MA, Largo R, Santillana J, Sanchez-Pernaute O, Calvo E, et al. (2006) Long term NSAID treatment inhibits COX-2 synthesis in the knee synovial membrane of patients with osteoarthritis: Differential proinflammatory cytokine profile between celecoxib and aceclofenac. *Ann Rheum Dis* 65: 998-1005.
133. Kingsbury SR, Tharmanathan P, Arden NK, Batley M, Birrell F, et al. (2015) Pain reduction with oral methotrexate in knee osteoarthritis, a pragmatic phase iii trial of treatment effectiveness (PROMOTE): Study protocol for a randomized controlled trial. *Trials* 16: 77.

Supplementary File

Methods

Search strategy

The literature search was performed on March 8, 2017 using the MedLine Advanced (PubMed) and the Embase Advanced search engines. The following keywords and their synonyms were used: Osteoarthritis, synovium or synovitis and macrophage. These keywords were used for searching titles and abstracts and were introduced

Supplementary Table 1: Overview of the terms used in both MedLine and Embase search.

Keyword	Synonyms
Osteoarthritis	– Osteoarthritis – Osteoarthrosis – Degenerative Joint Disease
Synovium	– Synovial Membrane – Synovial Tissue
Synovitis	– Synovial inflammation
Macrophage	– Macrophage – Monocyte – Histiocyte
Knee	

in the Medical Subject Heading (MeSH) vocabulary thesaurus system or Embase controlled vocabulary (EMTREE) (Supplementary Table 1 and Supplementary Table 2). A total of 389 records were retrieved after duplicates were removed. When retrieved articles fulfilled each of the inclusion criteria based on their title and abstract (N = 97) or when not, all criteria could be judged from title and abstract (N = 27), they were fully read for deciding on definite eligibility. Potentially relevant citations in fully read articles that were not among the MedLine and Embase search results were retrieved and manually included when they fulfilled each of the inclusion criteria (N = 6). As a result, a total of 104 articles was included in this study.

Inclusion and exclusion criteria

Supplementary Figure 1 depicts a flow diagram of the stepwise selection of the records. The search was limited to English research articles. Studies presenting non-original data, such as reviews, were excluded. Conference presentation abstracts were included if they contained retrievable data. Furthermore, experimental studies in non-human subjects and *in vitro* experiments were included.

Supplementary Table 2: Overview of the search strategies used.

Search engine	Search strategy	Result (number of records)
MedLine	(osteoarthritis* OR osteoarthrosis OR degenerative joint diseases*) AND knee AND (synovi* OR synovitis OR synovium) AND (macrophag* OR monocy* OR histiocy*) NOT review [Publication Type] AND English[Language]	176
Embase	osteoarthritis* OR osteoarthros* OR 'degenerative joint diseases*' AND (synovi* OR synovitis) AND (macrophag* OR monocy* OR histiocy*) NOT review:it AND [english]/lim AND knee	389

